Conservation Status of Eurasian Golden Plover (*Pluvialis apricaria*) and Song Thrush (*Turdus philomelos*)

2026 update

Wild Birds Regulation Unit

August 2026

(Link to the EEA Article 12 [2019–2024] Interactive Dashboard updated October 2026)

Contents

Executive Summary ii

Key to Conservation Status Codes iii

1. Quantitative Methodology for Reference Population Trends 1

2. Conservation Status of Eurasian Golden Plover (*Pluvialis apricaria*) 3

3. Conservation Status of Song Thrush (*Turdus philomelos*) 29

4. Conclusion 45

Executive Summary

This report provides an update on the conservation status of the Golden Plover (*Pluvialis apricaria*) and the Song Thrush (*Turdus philomelos*) for the Malta Ornis Committee.

The conservation status of both species is reviewed based on current scientific literature, which also provides details on population size, mortality rates, and migratory routes. The report integrates key findings from a satellite tagging project conducted between February 2022 and December 2025[1]. This study aimed to identify the breeding grounds and migratory patterns of these two species, with a focus on Malta's reference populations.

Key findings from the satellite tagging project

The satellite tagging project provided crucial data that corroborates and expands upon existing scientific literature.

- **Golden Plover**: The project yielded comprehensive data from 17 specimens. The study found that these birds wintered in Tunisia before migrating north and spent the breeding season in Sweden (n=3), Finland (n=3), Norway (n=4), and Russia (n=7). Overall, 35.3% of the tracked Golden Plovers bred within the territory of the European Union.

- **Song Thrush**: The study successfully tracked 21 specimens, with all birds wintering in Malta. Seven specimens provided conclusive evidence of their breeding locations in Romania (n=2), Bulgaria (n=2), Slovakia (n=1), Hungary (n=1), and Ukraine (n=1), confirming that 85.7% of the tracked Song Thrushes spent the breeding season within the territory of the European Union. This component also confirmed a distinct SW-NE pre-nuptial migration route.

Species-specific conservation assessment

Literature review, including the latest available ornithological publications and scientific evidence, confirms that Golden Plovers migrating over Malta are the subspecies *Pluvialis apricaria altifrons* from the Northeast European (West Continental) population. The GPS tracking data corroborated this finding and provided novel data showing that the Siberian population of the Eurasian Golden Plover also passes through Malta during both the pre-nuptial and post-nuptial migration periods.

The report includes a review of the latest population trend estimates for both species, based on the current Birds Directive Article 12 report (2019–2024) and the latest Pan-European Common Birds Monitoring Scheme (PECBMS) wild bird indicators. It also determines the applicable annual mortality rate for both species in accordance with the European Commission's Guide to Sustainable Hunting under the Birds Directive[2].

Data from the current Birds Directive Article 12 report (2019–2024) shows a short-term decreasing trend for Malta's EU reference population of *Pluvialis apricaria altifrons* (best value -10.97% based on combined Finnish and Swedish population data, with a minimum of -30% and maximum of +6% derived from Finnish data only). The species was reported as having a stable long-term trend (best value +10.96%, minimum -10.94%, maximum 35.74%).

Malta's EU reference population of *Turdus philomelos* shows a short-term increasing trend (best value +76.13%, minimum +27.99%, maximum +76.96%), while its long-term trend is unknown due to missing data from three of the four constituent Member States.

[1] Wild Birds Regulation Unit (WBRU), *Satellite Tagging of Song Thrushes* (Turdus philomelos) *and Eurasian Golden Plovers* (Pluvialis apricaria altifrons)*: Results from the 2022 – 2025 Research Period*, Final Report (2026). Available online: https://wbru.gov.mt/wp-content/uploads/2026/09/Final_Report_GP-ST-satellite-tagging-2022-2025.pdf.

[2] http://ec.europa.eu/environment/nature/conservation/wildbirds/hunting/docs/hunting_guide_en.pdf.

Category	European species of global conservation concern	Conservation status in Europe	Global population or range concentrated in Europe
SPEC 1	Yes	–	–
SPEC 2	No	Unfavourable	Yes
SPEC 3	No	Unfavourable	No
Non-SPEC[E]	No	Favourable	Yes
Non-SPEC	No	Favourable	No

Source: BirdLife International (2004: xiii)

Categories of Species of European Conservation Concern (SPECs) and Non-SPECs

Each species is initially assessed against the IUCN Red List Criteria (IUCN 2001) at a European level, and then against the additional criteria derived mainly from Birds in Europe I (Tucker and Heath 1994). All population size thresholds refer to minimum population estimates. In descending order of threat, a species is evaluated as:	
Critically Endangered (CR)	if its European population meets any of the IUCN Red List Criteria (A to E) for Critically Endangered. Such species have an Unfavourable conservation status in Europe because they are considered to be facing an extremely high risk of extinction in the wild (IUCN 2001).
Endangered (EN)	if its European population meets any of the IUCN Red List Criteria (A to E) for Endangered. Such species have an Unfavourable conservation status in Europe because they are considered to be facing a very high risk of extinction in the wild (IUCN 2001).
Vulnerable (V)	if its European population meets any of the IUCN Red List Criteria (A to E) for Vulnerable. Such species have an unfavourable conservation status in Europe because they are considered to be facing a high risk of extinction in the wild (IUCN 2001).
Declining (D)	if its European population does not meet any IUCN Red List Criteria, but declined by more than 10% over 10 years (i.e. 1990–2000) or three generations, whichever is longer. Such species have an Unfavourable conservation status in Europe because they are unable to maintain their populations and/or natural ranges in the long-term. [Birds in Europe I classified species as SPECs if the size of their population or range declined between 1970–1990 by **20%** or more in 33–65% of the population (or by 50% or more in 12–24% of the population). Given the shorter time period covered by Birds in Europe II, an overall decline exceeding **10%** is comparable with this approach.]
Rare (R)	if its European population does not meet any IUCN Red List Criteria and is not Declining, but numbers fewer than 10,000 breeding pairs (*or* 20,000 breeding individuals *or* 40,000 wintering individuals), and is not marginal to a larger non-European population. Such species have an Unfavourable conservation status in Europe because the small size of their population renders them more susceptible to accelerated declines as a result of: • break-up of social structure; • loss of genetic diversity; • large-scale population fluctuations and catastrophic chance events; • existing or potential exploitation, persecution or disturbance by humans.
Depleted (H)	if its European population does not meet any IUCN Red List Criteria and is not Rare or Declining, but has not yet recovered from a moderate or large decline suffered during 1970–1990, which led to its classification as Endangered, Vulnerable or Declining in Birds

	in Europe I. Such species have an Unfavourable conservation status in Europe because they have already undergone a population decline of the type that various directives, conventions and agreements intend to prevent, and have not yet recovered.
Localised (L)	if its European population does not meet any IUCN Red List Criteria and is not Declining, Rare or Depleted, but is heavily concentrated, with more that 90% of the European population occurring at 10 or fewer sites (as listed in Heath and Evans 2000). Such species have an Unfavourable conservation status in Europe because their dependence on a small number of sites renders them more susceptible to accelerated declines as a result of: large-scale population fluctuations and catastrophic chance events; existing or potential exploitation, persecution and disturbance by humans.
Secure (S)	if its European population does not meet any of the criteria listed above. Such species have a Favourable conservation status in Europe.
In addition, a species is considered to be:	
Data Deficient (DD)	if there is inadequate information to make a direct, or indirect, assessment of its risk of extinction based on its distribution and/or population status. A species in this category may be well studied, and its biology well known, but appropriate data on its abundance and/or distribution in Europe are lacking. Data Deficient is therefore not a category of threat (IUCN 2001).
Not evaluated (NE)	if its European population has not yet been evaluated against the criteria.
Source: BirdLife International (2004: 8)	

Species trends in Birds in Europe (2004)

'Worst case' trend scenario 1990–2000	1990–2000 trend category	Criteria met
>30% decline	Large decline	IUCN Red List Criteria
10–29% decline	Moderate decline	Declining
<10% decline and <10% increase	Stable	-
10–29% increase	Moderate increase	-
>30% increase	Large increase	-
Unknown (insufficient data)	Unknown	-


Source: BirdLife International (2004)

IUCN Categories

EXTINCT (EX)	A taxon is Extinct when there is no reasonable doubt that the last individual has died. A taxon is presumed Extinct when exhaustive surveys in known and/or expected habitat, at appropriate times (diurnal, seasonal, annual), throughout its historic range have failed to

	record an individual. Surveys should be over a time frame appropriate to the taxon's life cycle and life form.
EXTINCT IN THE WILD (EW)	A taxon is Extinct in the Wild when it is known only to survive in cultivation, in captivity or as a naturalized population (or populations) well outside the past range. A taxon is presumed Extinct in the Wild when exhaustive surveys in known and/or expected habitat, at appropriate times (diurnal, seasonal, annual), throughout its historic range have failed to record an individual. Surveys should be over a time frame appropriate to the taxon's life cycle and life form.
CRITICALLY ENDANGERED (CR)	A taxon is Critically Endangered when the best available evidence indicates that it meets any of the criteria A to E for Critically Endangered (see Section V), and it is therefore considered to be facing an extremely high risk of extinction in the wild.
ENDANGERED (EN)	A taxon is Endangered when the best available evidence indicates that it meets any of the criteria A to E for Endangered (see Section V), and it is therefore considered to be facing a very high risk of extinction in the wild.
VULNERABLE (VU)	A taxon is Vulnerable when the best available evidence indicates that it meets any of the criteria A to E for Vulnerable (see Section V), and it is therefore considered to be facing a high risk of extinction in the wild.
NEAR THREATENED (NT)	A taxon is Near Threatened when it has been evaluated against the criteria but does not qualify for Critically Endangered, Endangered or Vulnerable now, but is close to qualifying for or is likely to qualify for a threatened category in the near future.
LEAST CONCERN (LC)	A taxon is Least Concern when it has been evaluated against the criteria and does not qualify for Critically Endangered, Endangered, Vulnerable or Near Threatened. Widespread and abundant taxa are included in this category.
DATA DEFICIENT (DD)	A taxon is Data Deficient when there is inadequate information to make a direct, or indirect, assessment of its risk of extinction based on its distribution and/or population status. A taxon in this category may be well studied, and its biology well known, but appropriate data on abundance and/or distribution are lacking. Data Deficient is therefore not a category of threat. Listing of taxa in this category indicates that more information is required and acknowledges the possibility that future research will show that threatened classification is appropriate. It is important to make positive use of whatever data are available. In many cases great care should be exercised in choosing between DD and a threatened status. If the range of a taxon is suspected to be relatively circumscribed, and a considerable period of time has elapsed since the last record of the taxon, threatened status may well be justified.
NOT EVALUATED (NE)	A taxon is Not Evaluated when it has not yet been evaluated against the criteria.

Structure of IUCN categories



Source: IUCN Red List Categories and Criteria Ver. 3.1 2nd edition. Available at: https://portals.iucn.org/library/efiles/documents/RL-2001-001-2nd.pdf

IUCN CRITERIA FOR CRITICALLY ENDANGERED, ENDANGERED AND VULNERABLE TAXA

CRITICALLY ENDANGERED (CR)	A taxon is Critically Endangered when the best available evidence indicates that it meets any of the following criteria (A to E), and it is therefore considered to be facing an extremely high risk of extinction in the wild: A. Reduction in population size based on any of the following: 1. An observed, estimated, inferred or suspected population size reduction of ≥90% over the last 10 years or three generations, whichever is the longer, where the causes of the reduction are clearly reversible AND understood AND ceased, based on (and specifying) any of the following: (a) direct observation (b) an index of abundance appropriate to the taxon (c) a decline in area of occupancy, extent of occurrence and/or quality of habitat (d) actual or potential levels of exploitation (e) the effects of introduced taxa, hybridization, pathogens, pollutants, competitors or parasites. 2. An observed, estimated, inferred or suspected population size reduction of ≥80% over the last 10 years or three generations, whichever is the longer, where the reduction or its causes may not have ceased OR may not be understood OR may not be reversible, based on (and specifying) any of (a) to (e) under A1. 3. A population size reduction of ≥80%, projected or suspected to be met within the next 10 years or three generations, whichever is the longer (up to a maximum of 100 years), based on (and specifying) any of (b) to (e) under A1. 4. An observed, estimated, inferred, projected or suspected population size reduction of ≥80% over any 10 year or three generation period, whichever is longer (up to a maximum of 100 years in the future), where the time period must include both the past and the future, and where the reduction or its causes may not have ceased OR may not be understood OR may not be reversible, based on (and specifying) any of (a) to (e) under A1. B. Geographic range in the form of either B1 (extent of occurrence) OR B2 (area of occupancy) OR both:

	1. Extent of occurrence estimated to be less than 100 km^2, and estimates indicating at least two of a-c: a. Severely fragmented or known to exist at only a single location. b. Continuing decline, observed, inferred or projected, in any of the following: (i) extent of occurrence (ii) area of occupancy (iii) area, extent and/or quality of habitat (iv) number of locations or subpopulations (v) number of mature individuals. c. Extreme fluctuations in any of the following: (i) extent of occurrence (ii) area of occupancy (iii) number of locations or subpopulations (iv) number of mature individuals. 2. Area of occupancy estimated to be less than 10 km^2, and estimate indicating at least two of a-c: a. Severely fragmented or known to exist at only a single location. b. Continuing decline, observed, inferred or projected, in any of the following: (i) extent of occurrence (ii) area of occupancy (iii) area, extent and/or quality of habitat (iv) number of locations or subpopulations (v) number of mature individuals. c. Extreme fluctuations in any of the following: (i) extent of occurrence (ii) area of occupancy (iii) number of locations or subpopulations (iv) number of mature individuals. C. Population size estimated to number fewer than 250 mature individuals and either: 1. An estimated continuing decline of at least 25% within three years or one generation, whichever is longer, (up to a maximum of 100 years in the future) OR 2. A continuing decline, observed, projected, or inferred, in numbers of mature individuals AND at least one of the following (a-b): . Population structure in the form of one of the following: (i) no subpopulation estimated to contain more than 50 mature individuals, OR (ii) at least 90% of mature individuals in one subpopulation. b. Extreme fluctuations in number of mature individuals. D. Population size estimated to number fewer than 50 mature individuals. E. Quantitative analysis showing the probability of extinction in the wild is at least 50% within 10 years or three generations, whichever is the longer (up to a maximum of 100 years).
ENDANGERED (EN)	A taxon is Endangered when the best available evidence indicates that it meets any of the following criteria (A to E), and it is therefore considered to be facing a very high risk of extinction in the wild: . Reduction in population size based on any of the following: 1. An observed, estimated, inferred or suspected population size reduction of ≥70% over the last 10 years or three generations, whichever is the longer, where the causes of the reduction are clearly reversible AND understood AND ceased, based on (and specifying) any of the following:

	(a) direct observation (b) an index of abundance appropriate to the taxon (c) a decline in area of occupancy, extent of occurrence and/or quality of habitat (d) actual or potential levels of exploitation (e) the effects of introduced taxa, hybridization, pathogens, pollutants, competitors or parasites. 2. An observed, estimated, inferred or suspected population size reduction of ≥50% over the last 10 years or three generations, whichever is the longer, where the reduction or its causes may not have ceased OR may not be understood OR may not be reversible, based on (and specifying) any of (a) to (e) under A1. 3. A population size reduction of ≥50%, projected or suspected to be met within the next 10 years or three generations, whichever is the longer (up to a maximum of 100 years), based on (and specifying) any of (b) to (e) under A1. 4. An observed, estimated, inferred, projected or suspected population size reduction of ≥50% over any 10 year or three generation period, whichever is longer (up to a maximum of 100 years in the future), where the time period must include both the past and the future, AND where the reduction or its causes may not have ceased OR may not be understood OR may not be reversible, based on (and specifying) any of (a) to (e) under A1. B. Geographic range in the form of either B1 (extent of occurrence) OR B2 (area of occupancy) OR both: 1. Extent of occurrence estimated to be less than 5,000 km^2, and estimates indicating at least two of a-c: a. Severely fragmented or known to exist at no more than five locations. b. Continuing decline, observed, inferred or projected, in any of the following: (i) extent of occurrence (ii) area of occupancy (iii) area, extent and/or quality of habitat (iv) number of locations or subpopulations (v) number of mature individuals. c. Extreme fluctuations in any of the following: (i) extent of occurrence (ii) area of occupancy (iii) number of locations or subpopulations (iv) number of mature individuals. 2. Area of occupancy estimated to be less than 500 km^2, and estimates indicating at least two of a-c: a. Severely fragmented or known to exist at no more than five locations. b. Continuing decline, observed, inferred or projected, in any of the following: (i) extent of occurrence (ii) area of occupancy (iii) area, extent and/or quality of habitat (iv) number of locations or subpopulations (v) number of mature individuals. c. Extreme fluctuations in any of the following: (i) extent of occurrence (ii) area of occupancy (iii) number of locations or subpopulations (iv) number of mature individuals. C. Population size estimated to number fewer than 2,500 mature individuals and either: 1. An estimated continuing decline of at least 20% within five years or two generations, whichever is longer, (up to a maximum of 100 years in the future) OR 2. A continuing decline, observed, projected, or inferred, in numbers of mature individuals AND at least one of the following (a-b):

	a. Population structure in the form of one of the following: (i) no subpopulation estimated to contain more than 250 mature individuals, OR (ii) at least 95% of mature individuals in one subpopulation. b. Extreme fluctuations in number of mature individuals. D. Population size estimated to number fewer than 250 mature individuals. E. Quantitative analysis showing the probability of extinction in the wild is at least 20% within 20 years or five generations, whichever is the longer (up to a maximum of 100 years).
VULNERABLE (VU)	A taxon is Vulnerable when the best available evidence indicates that it meets any of the following criteria (A to E), and it is therefore considered to be facing a high risk of extinction in the wild: . Reduction in population size based on any of the following: 1. An observed, estimated, inferred or suspected population size reduction of ≥50% over the last 10 years or three generations, whichever is the longer, where the causes of the reduction are clearly reversible AND understood AND ceased, based on (and specifying) any of the following: (a) direct observation (b) an index of abundance appropriate to the taxon (c) a decline in area of occupancy, extent of occurrence and/or quality of habitat (d) actual or potential levels of exploitation (e) the effects of introduced taxa, hybridization, pathogens, pollutants, competitors or parasites. 2. An observed, estimated, inferred or suspected population size reduction of ≥30% over the last 10 years or three generations, whichever is the longer, where the reduction or its causes may not have ceased OR may not be understood OR may not be reversible, based on (and specifying) any of (a) to (e) under A1. 3. A population size reduction of ≥30% projected or suspected to be met within the next 10 years or three generations, whichever is the longer (up to a maximum of 100 years), based on (and specifying) any of (b) to (e) under A1. 4. An observed, estimated, inferred, projected or suspected population size reduction of ≥30% over any 10 year or three generation period, whichever is longer (up to a maximum of 100 years in the future), where the time period must include both the past and the future, AND where the reduction or its causes may not have ceased OR may not be understood OR may not be reversible, based on (and specifying) any of (a) to (e) under A1. B. Geographic range in the form of either B1 (extent of occurrence) OR B2 (area of occupancy) OR both: 1. Extent of occurrence estimated to be less than 20,000 km^2, and estimates indicating at least two of a-c: a. Severely fragmented or known to exist at no more than 10 locations. b. Continuing decline, observed, inferred or projected, in any of the following: (i) extent of occurrence (ii) area of occupancy (iii) area, extent and/or quality of habitat (iv) number of locations or subpopulations (v) number of mature individuals. c. Extreme fluctuations in any of the following: (i) extent of occurrence (ii) area of occupancy (iii) number of locations or subpopulations (iv) number of mature individuals. 2. Area of occupancy estimated to be less than 2,000 km^2, and estimates indicating at least two of a-c: a. Severely fragmented or known to exist at no more than 10 locations.

	b. Continuing decline, observed, inferred or projected, in any of the following: (i) extent of occurrence (ii) area of occupancy (iii) area, extent and/or quality of habitat (iv) number of locations or subpopulations (v) number of mature individuals. c. Extreme fluctuations in any of the following: (i) extent of occurrence (ii) area of occupancy (iii) number of locations or subpopulations (iv) number of mature individuals. C. Population size estimated to number fewer than 10,000 mature individuals and either: 1. An estimated continuing decline of at least 10% within 10 years or three generations, whichever is longer, (up to a maximum of 100 years in the future) OR 2. A continuing decline, observed, projected, or inferred, in numbers of mature individuals AND at least one of the following (a-b): a. Population structure in the form of one of the following: (i) no subpopulation estimated to contain more than 1,000 mature individuals, OR (ii) all mature individuals in one subpopulation. b. Extreme fluctuations in number of mature individuals. D. Population very small or restricted in the form of either of the following: 1. Population size estimated to number fewer than 1,000 mature individuals. 2. Population with a very restricted area of occupancy (typically less than 20 km^2) or number of locations (typically five or fewer) such that it is prone to the effects of human activities or stochastic events within a very short time period in an uncertain future, and is thus capable of becoming Critically Endangered or even Extinct in a very short time period. E. Quantitative analysis showing the probability of extinction in the wild is at least 10% within 100 years.

1. Quantitative Methodology for Reference Population Trends

Overview and Analytical Purpose

To assess the conservation status of Malta's European Union (EU) reference population of Eurasian Golden Plover (*Pluvialis apricaria altifrons*) and Song Thrush (*Turdus philomelos*), quantitative assessment protocols integrate satellite tracking movement data[3] with Article 12 Birds Directive reporting data (2019–2024).

Employing spatial demography (spatial weighting), this approach accounts for species-specific migratory connectivity by scaling reference populations according to verified breeding grounds. To evaluate trend magnitudes and net population changes, calculations anchor directly to historical baseline reconstruction. Reconstructing starting populations ($P_{\text{ref, baseline}}$) measures the net gain or loss of breeding pairs relative to their original baseline.

Direct cross-references corresponding to the datasets for Eurasian Golden Plover and Song Thrush are explicitly indicated throughout this section.

Derivation of Weighted Reference Populations

The reference population is determined by scaling breeding pair figures according to satellite tracking spatial distribution:

$$P_{\text{ref, current}} = \sum (P_{\text{min}} \times \alpha)$$

Where:

- $P_{\text{ref, current}}$ is Malta's weighted EU reference population in minimum breeding pairs (corresponds to column header *Malta's EU reference population (Min. Pairs)*).
- P_{min} is the minimum breeding population reported by a given EU Member State (corresponds to column header *Min. Breeding Pairs*).
- α is the weighted reference proportion of satellite-tracked individuals that spent the breeding season in that specific Member State (corresponds to column header *Weighted Share of Tracked Birds*).

Baseline Reconstruction and Absolute Change Calculation

Population trend magnitudes reported under Article 12 reporting (T) reflect proportional changes relative to a historical baseline ($P_{\text{ref, baseline}}$). To compute the actual change in breeding pairs (ΔP) directly from current reference population figures, the historical starting baseline is reconstructed using the inverse rate equation:

$$P_{\text{ref, baseline}} = \frac{P_{\text{ref, current}}}{1 + T}$$

Where:

- $P_{\text{ref, baseline}}$ is Malta's reconstructed weighted EU reference baseline population in minimum breeding pairs (derived parameter based on *Malta's EU reference population (Min. Pairs)*).

[3] Wild Birds Regulation Unit (WBRU), *Satellite Tagging of Song Thrushes* (Turdus philomelos) *and Eurasian Golden Plovers* (Pluvialis apricaria altifrons)*: Results from the 2022 – 2025 Research Period*, Final Report (2026). Available online: https://wbru.gov.mt/wp-content/uploads/2026/09/Final_Report_GP-ST-satellite-tagging-2022-2025.pdf.

- $P_{\text{ref, current}}$ is Malta's current weighted EU reference population in minimum breeding pairs (corresponds to column header *Malta's EU reference population (Min. Pairs)*).

- T is the reported percentage trend magnitude expressed as a decimal (corresponds to column header *Trend Magnitude (%)*; e.g., a -15% decrease is expressed as -0.15; a $+5\%$ increase is expressed as $+0.05$).

Once the weighted historical baseline is established, the net absolute change in breeding pairs (ΔP) relative to that starting point is calculated by subtracting the baseline from the current reference population:

$$\Delta P = P_{\text{ref, current}} - P_{\text{ref, baseline}}$$

Where:

- ΔP is the net absolute gain or loss of breeding pairs relative to the weighted reference baseline (corresponds to the section header *Calculated Change in Breeding Pairs (Relative to Historical Baseline)*).

This two-step calculation guarantees complete mathematical consistency whether population trends are positive, negative, or stable.

Data Bound Evaluation and Handling of Missing Values

To maintain analytical rigour, dataset parameters and missing values are treated under strict operational rules:

- **Reported Trend Ranges (Min/Max):** Where Member States report a range for population trends (e.g., $T_{\text{min}} = -10\%$; $T_{\text{max}} = +10\%$), absolute changes are calculated independently at both boundary parameters (corresponds to sub-columns *Min* and *Max*) to capture full scientific uncertainty.

- **Unreported Data (N/A):** Where a specific parameter (e.g., *Best Value*) is not reported by a country, it is treated as an intentional omission (N/A). Non-reporting countries are excluded exclusively from that parameter's subset calculation rather than assigned a value of zero.

- **Confirmed Zero Values ($\boldsymbol{T = 0\%}$):** Explicitly reported 0% trend magnitudes represent confirmed, reported data points. These values are retained in all calculations to maintain the Member State's full contribution to baseline denominators.

- **Overall Aggregated Trend Percentage ($\boldsymbol{T_{\text{agg}}}$):** Corresponds directly to the *Overall Population Change (%)* summary row. This parameter quantifies the overall proportional change of an active subset by dividing the net sum of absolute population changes ($\sum \Delta P$) across all contributing Member States by the total reconstructed baseline population ($\sum P_{\text{ref, baseline}}$) of the contributing countries in that calculation:

$$T_{\text{agg}} = \frac{\sum \Delta P}{\sum P_{\text{ref, baseline}}} \times 100$$

2. Conservation Status of Eurasian Golden Plover (*Pluvialis apricaria*)

The Golden Plover (*Pluvialis apricaria s.l.*) is a medium-sized Palearctic wader which breeds in continental arctic, arctic alpine and boreal tundra, but secondarily on temperate oceanic unenclosed upland moors and peat-land. In winter, the Golden Plover occupies harvest fields, stubbles, close grazed pastures, fallows and other farmland of open character including flood-lands. Its breeding range extends from Iceland to Russia, and its wintering quarters stretch from Morocco to Asia Minor including South Europe (Geroudet 1983; Cramp & Simmons 1983).

The Golden Plover is a huntable species listed under Annex II/B of the Birds Directive. According to the EU Management Plan for the Golden Plover[4], this species is hunted in at least four EU Member States, including Portugal, Ireland, France and Malta. According to the Management Plan, over 62,000 Golden Plovers were known to have been bagged annually in France and over 32,000 in Portugal[5].

IUCN Red List Category – Least Concern

This species has an extremely large range and hence does not approach the thresholds for Vulnerable under the range size criterion (extent of occurrence under 20,000 km² combined with a declining or fluctuating range size, habitat extent/quality, or population size and a small number of locations or severe fragmentation). The population size is extremely large and hence does not approach the thresholds for Vulnerable under the population size criterion (under 10,000 mature individuals with a continuing decline estimated to be over 10% in ten years or three generations, or with a specified population structure). The population trend appears to be stable, and hence the species does not approach the thresholds for Vulnerable under the population trend criterion (over 30% decline over ten years or three generations). For these reasons this species is evaluated by BirdLife International (2026) and IUCN[6] as **Least Concern**.

Sub-specific assessment – a historical review of trends and geographical populations

Cramp and Simmons (1983)[7] maintain that the Golden Plover includes two sub-species ***Pluvialis apricaria apricaria*** (nominate) breeding from Ireland and Britain to the Baltic States and ***Pluvialis apricaria altifrons*** breeding at higher altitudes from Iceland to north-central Siberia. Delany *et. al*. (2007)[8] indicate that the *altifrons* population is divided into three sub-populations, of which the Northeast European (West Continental) is the one that migrates over Malta (breeds from North Norway to Russia-east 70 degrees) and winters in west and south continental Europe, east Britain and north-west Africa (Morocco to Tunisia).

The Agreement on the Conservation of African–Eurasian Migratory Waterbirds (AEWA) Status Report indicates that the *Pluvialis apricaria altifrons* breeds widely at high altitudes in Western Eurasia from Iceland east to 100 degrees east in Western Siberia, and winters south to North Africa (occasionally to Senegal) and the South Caspian. *Pluvialis apricaria apricaria* breeds at more westerly latitudes in Britain, Ireland, Denmark and Germany, and is a short-distance migrant, most birds wintering fairly close to their breeding areas (Scott, 2002).

---

[4] European Union Management Plan 2009–2011 Golden Plover *Pluvialis apricaria*. Available at: https://ec.europa.eu/environment/nature/conservation/wildbirds/hunting/docs/Golden%20Plover%20EU_MP.pdf.

[5] EC 2009: *European Union Management Plan for Golden Plover 2009–2011*, Table 4, page 17.

[6] http://www.iucnredlist.org/details/22693727/0. Downloaded on 31/08/2026.

[7] Cramp, S., cf. ed. (1983). The Birds of the Western Palearctic. Vol. III 'Waders to Gulls. (Pluvialis apricaria Golden Plover pp 201-216). Oxford University Press.

[8] Delany, S, Scott, D, Dodman T, and Stroud, D (eds) (2009) *2009 Atlas of Wader Populations in Africa and Western Eurasia.* Wetlands International, Wageningen, The Netherlands.

Considerable variation exists in most populations, and many authors consider the species to be monotypic. Four main breeding populations can be identified: (1) Icelandic and Faeroese breeders (*altifrons*), wintering mainly in Ireland, with smaller numbers in western Britain, France and Iberia, and a few to Northwest Africa; (2) a population of *altifrons* breeding from Northern Norway east to about 70 degrees east and wintering mainly in western and southern continental Europe and Northwest Africa (Morocco and Tunisia) [Malta's EU reference population]; (3) population of *altifrons* breeding in Western Siberia east to 100 degrees east and wintering in the Caspian region band possibly Asia Minor and Eastern Mediterranean; and (4) southern breeders (*apricaria*) breeding in Britain, Ireland, Denmark (extirpated?) and Germany, and wintering in Northwest Europe (Table 1).

Table 1. Countries of origin of *Pluvialis apricaria altifrons* and *Pluvialis apricaria apricaria*

Population	***altifrons* (Iceland & Faeroes)**	***altifrons* (NW Europe)**	***altifrons* (N–C Siberia)**	***apricaria* (UK/IR/DK/D)**
Population size	☺	☺	☹☹	☹
Breeding location	☺☺	☺☺	😐	☺☺

Source: Adapted from Gillings, S (2005) Table 1: An assessment of the level of knowledge for developing a flyway conservation strategy for each breeding population of Eurasian Golden Plover (after Davidson *et al*. 1998). ☺☺ = very good; ☺ = good; = 😐 adequate; ☹ = poor, ☹☹ = very poor. International workshop on passage and wintering Eurasian Golden Plovers (p. 6).

AEWA and its Resolution 4.11 (see http://tinyuri.com/aewares411) also provides the distribution for *Pluvialis apricaria apricaria* as "*Britain, Ireland, Denmark, Germany and the Baltic*", thereby excluding in a definitive way the Central Mediterranean region. The same applies for the 'European Management Plan for the Golden Plover, 2009–2011' (see http://tinyuri.com/gp-eump) which indicates that "*The nominate southern subspecies Pluvialis apricaria apricaria nests in Ireland, Great Britain, Denmark, Germany, Latvia, Lithuania and Estonia. This subspecies winters in North-West, from Ireland to South Britain, France and Iberia*".

In this context, it should be noted that the sub-species that migrates over Malta in autumn is *Pluvialis apricaria altifrons* as can also be attested by Delany *et. al*. (2007). The authors maintain that "there is growing evidence that numbers of *altifrons* wintering in Europe are increasing" (Delany et. al., 2007: 183). The publication also indicates that the nominate *Pluvialis apricaria apricaria* has a more restricted migratory distribution that excludes the Maltese Islands and nearby central regions in the Mediterranean. The European Union Management Plan (EC 2009) specifies that:

"*The Northern P. a. altifrons subspecies is generally split into three populations based on their different flyways that overlap in winter. They are defined as:*

- *Icelandic (or East Atlantic): breeds in Iceland, the Faeroe Islands and Greenland and winters in Ireland, West Britain, France, West Iberia and North-West Africa.*
- *Northeast European (or West Continental): breeds in North Norway and Russia (east to 70 degrees east) and winters in West and south continental Europe, East Britain and North-West Africa (Morocco and Tunisia).*
- *North Central Siberia population, breeds in North Central Siberia (east to 100 degrees east) and winters in the Caspian Region, Asia Minor and the Eastern Mediterranean."*

According to Delany *et al*. (2007) *Pluvialis apricaria apricaria* is declining whereas *Pluvialis apricaria altifrons* is more or less stable (see also page 7, second paragraph, of *The European Union Management Plan 2009-2011 for Golden Plover*). On the basis of current available literature, therefore, the population of the sub-species with a historical decline is the *P. a. apricaria*, whereas the other populations / groups (*P. apricaria altifrons*) found in Europe are stable or increasing.

According to the EU Management Plan for the Golden Plover, approximately 50% of Golden Plover (*P. apricaria altifrons*) European population breeds in Iceland with a total of >300,000 breeding pairs (Thorup 2006, 930,000 individuals, Wetlands International 2006). Trends in this population are not well known, whereas the relatively small Faeroese population is declining. Approximately half of this population is found in Norway, 22% in Finland, 17% in Sweden and 9% in Russia with a total ranging from 217,000 to 362,000 breeding pairs. This population is considered stable (Wetlands International 2006). According to AEWA, while the *P. apricaria apricaria* is declining, *P. apricaria altifrons* is considered as stable and possibly increasing.

It should thus be noted that this moderate decline is related to the nominate *Pluvialis apricaria apricaria* population which does not pass through Malta (Figures 1 and 2). Golden Plovers that migrate over the Maltese Islands form part of the Northeast European (or West Continental) population (*P. apricaria altifrons*) which has a **minimum population of 172,089 breeding pairs at EU level** (Finland and Sweden only) – see Table 5.




Figure 1. Outline of the timing and geography of the southward migration of Golden Plovers. Arrows indicate the most important migration routes, and size of arrow is an indication of migration intensity. Source: Jukema *et al.* 2001. In: Gillings, S (2005) International workshop on passage and wintering Eurasian Golden Plovers (p. 5).




Figure 2. Migratory flyways of European (Eurasian) Golden Plover *Pluvialis apricaria*. Source: Wetlands International (2009) *An Atlas of Wader Populations in Africa and Western Eurasia* (p. 180). Note orange boundary delineating the restricted migration pattern of nominate *Pluvialis apricaria apricaria*.

Article 12 reports (2019–2024)

This update includes the current population trends and indices of Malta's EU reference population as reported by Member States in their Article 12 report for period 2019–2024[9]. The Article 12 (2019–2024) dashboard containing Member State data files has recently been made public, accessible at https://nature-art12.eionet.europa.eu/article12/summary. However, the overall EU population status for each species is still pending, awaiting the subsequent release of national summaries, final datasets, and the "State of Nature" report by December 2026.

During the previous reporting cycle (2013–2018), the Eurasian Golden Plover was reported as having a breeding population size of 207,000–338,000 pairs within the territory of the European Union (previously reported at 244,000–355,000 pairs during the 2008–2012 reporting cycle) and a breeding range size of 533,000km² (previously reported at 661,000km² during the 2008–2012 reporting cycle). During the 2013–2018 reporting cycle, the breeding population trend in the EU was reported as stable in both the short-term and long-term (EEA, 2020)[10].

Pan-European Common Birds Monitoring Scheme (2025 update)

It should be noted that since 2021, Estonia has been included in the "List of Countries" contributing to the latest European Bird Census Council (EBCC / PECBMS) updates, apart from Norway, **Sweden** and **Finland** (the latter two constituting Malta's EU reference population). According to the EU Management Plan for this species, Estonia, which sustains a breeding population of 2,500–3,400 pairs (EEA, 2020) does not form part of the Northeast European population of *Pluvialis apricaria altifrons* (Malta's EU reference population)[11].

According to the latest dataset available from the European Bird Census Council (EBCC, 2025)[12] at the time the conservation status of the Golden Plover (*Pluvialis apricaria*) was updated (August 2026), **there was a decrease in the combined breeding population of the contributing countries** (Norway, Sweden, Finland and Estonia) both in the short-term and long-term trends. When compared with the previous EBCC update, the decrease in the short-term trend [ten-year slope] (2015–2024) was from -11% to -16% (**an additional -5% decrease**) and from -31% to -41% (**an additional -10% decrease**) in the long-term trend (1980–2024)[13]. The current EBCC assessment considers *Pluvialis apricaria* as **Stable**.

Figure 3 is an extract from the EBCC 2025 update, which confirms that the population trend category for *Pluvialis apricaria* remained **Stable** with "no significant increase or decline, and most probable trends are less than 5% per year".

---

[9] Available at: https://nature-art12.eionet.europa.eu/article12/summary.

[10] European Environment Agency (2020) Article 12 reports for the 2013-2018 reporting period (EU). Available at: https://nature-art12.eionet.europa.eu/article12/summary?period=3&subject=Pluvialis+apricaria&reported_name=.

[11] *The nominate southern subspecies* P. a. apricaria *nests in Ireland, Great Britain, Denmark, Germany, Latvia, Lithuania and Estonia. This subspecies winters in North-West Europe, form Ireland to South Britain, France and Iberia.* Source: EU Management Plan – Golden Plover (p.8).

[12] *Trends of common birds in Europe, 2025 update*. Available at: https://pecbms.info/trends-and-indicators/species-trends/species/pluvialis-apricaria/.

[13] Comparison between the previous and current EBCC updates is for information purposes only. The EBCC updates provide composite data (combined data of all contributing countries) on the latest population trends at pan-European level, not at EU level—it includes data pertaining to countries that are not part of Malta's EU reference population. Moreover, since the EBCC updates do not provide segregated data on population levels and trends for each contributing country, it is not possible to use EBCC's composite data, even if combined at EU level only, to update the assessment of the species at Malta's EU reference population level. The data at EU level that is segregated by country, and hence available to the Maltese authorities to carry out an assessment of the status of the species at reference population level, is the one published by the European Environment Agency as part of the Birds Directive Article 12 reporting obligations.




Figure 3. Long-term trend classification for *Pluvialis apricaria*. Source: EBCC (2025 update). Available at: https://pecbms.info/trends-and-indicators/species-trends/species/pluvialis-apricaria/.

List of Countries:

Estonia	1984 – 2024
Finland	1981 – 2024
Norway	2007 – 2024
Sweden	1998 – 2024

The Golden Plover does not breed in Malta (neither in the wild, nor in captivity). Table 2 lists the live-capturing data for Golden Plover in Malta and Gozo from 2006 to December 2024.

Table 2. Live-capturing data for Golden Plover (2006–04 Jan 2026)

Species	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	Total	Mean (20-year average)
Golden Plover	319	44	503[1]	414	1,769	52	25[2]	158[3]	476[4]	321[5]	297[5]	167[5]	233[5]	588[6]	797[7]	688[8]	727[9]	700[10]	700[11]	700[12]	9,678	484

Data sources:

2006–2012: *Carnet de Chasse* data, Malta Environment and Planning Authority.
2013–2025: Game Reporting System, Wild Birds Regulation Unit.

[1] Jan 2008 (n=16) + Oct–Dec 2008 (n=487)
[2] The 2012 figure represents the SMS data for the period 20th October to 31st December 2012 (i.e. excluding the period 1st January to 10th January 2013, n=8).
[3] The 2013 figure represents the SMS data for the period 20th October to 31st December 2013 (i.e. excluding the period 1st January to 10th January 2014, n=58).
[4] The 2014 figure represents the SMS data for the period 1st January to 10th January 2014 (n=58) and the game reporting data for the period 20th October to 31st December 2014 (n=418).
[5] Game reporting data for the period 20th October to 31st December of the corresponding year.
[6] Game reporting data for the period 1st November to 31st December of the corresponding year.
[7] The 2020 figure represents the game reporting data for the period 1st January to 10th January 2020 (n=112) and the game reporting data for the period 1st November to 31st December 2020 (n=685).
[8] The 2021 figure represents the game reporting data for the period 1st January to 4th January 2021 (n=15) [season closed on 4th January 2021 as soon as the national bag limit of 700 Golden Plovers was reached] and the game reporting data for the period 1st November to 31st December 2021 (n=673).
[9] The 2022 figure represents the game reporting data for the period 1st January to 3rd January 2022 (n=27) [season closed on 3rd January 2022 as soon as the national bag limit of 700 Golden Plovers was reached] and the game reporting data for the period 1st November to 30th December 2022 (n=700). The 01 November 2022 – 10 January 2023 live-capturing season for Golden Plover was closed on 30th December 2022 as soon as the national bag limit of 700 Golden Plovers was reached.
[10] The 2023 figure represents the game reporting data for the period 1st November to 23rd December 2023 (n=700). The 01 November 2023 – 10 January 2024 live-capturing season for Golden Plover was closed on 23rd December 2023 as soon as the national bag limit of 700 Golden Plovers was reached.
[11] The 2024 figure represents the game reporting data for the period 1st November to 27th December 2024 (n=700). The 01 November 2024 – 10 January 2025 live-capturing season for Golden Plover was closed on 27th December 2024 as soon as the national bag limit of 700 Golden Plovers was reached.
[12] The 2025 figure represents the game reporting data for the period 1st November 2025 to 4th January 2026 (n=700). The 01 November 2025 – 10 January 2026 live-capturing season for Golden Plover was closed on 4th January 2026 as soon as the national bag limit of 700 Golden Plovers was reached.

Reference population

The "small numbers" requirement, within the meaning of Article 9(1)(c) of the Birds Directive should be understood as "any sample of less than 1% of the total annual mortality of the population in question (average value) for those species which are not to be hunted and **a sample in the order of 1% for those species which may be hunted**. 'Population in question' is to be understood, with regard to migratory species, as the population of those regions from which come the main contingents passing through the region to which the derogation applies during its period of application". Given that the Golden Plover qualifies as a huntable species under the Birds Directive (MT: Annex IIB), the "*small numbers*" requirement is understood as "*a sample in the order of 1% of the total annual mortality of the population in question*".

As explained in the previous sections of this report, on the basis of the latest available scientific literature and satellite tracking data, the reference population of the Golden Plover for Malta forms part of the West Continental (Northeast European) population (*P. apricaria altifrons*). According to the EU Management Plan (2009–2011) for this species, this population is distributed as follows:

- Norway (c. 50%)
- Finland (22%)
- Sweden (17%)
- Russia (9%)

Malta's EU reference population of the Golden Plover is Finland and Sweden.

Satellite tracking data

Although the flyways of Golden Plovers are mapped in established scientific literature (see Figures 1 and 2 above), as requested by the services of the European Commission in 2019, the Maltese authorities initiated a research project to assess the migratory routes of this species passing through Malta using satellite tracking devices, hereinafter referred to as satellite tags. The location of tagged specimens was monitored by the Maltese authorities using Argos satellite data repository to which a Movebank account was linked. The Movebank account showing all GPS fixes was converted to a publicly visible research study in 2021. This conservation status report incorporates the final findings of the study[14].



The project yielded comprehensive data from 17 Golden Plovers. The study found that these birds wintered in Tunisia before migrating north and spent the breeding season in Sweden (n=3), Finland (n=3), Norway (n=4), and Russia (n=7). Overall, 35.3% of the tracked Golden Plovers bred within the EU.



The GPS fixes for the satellite-tagged Golden Plovers are detailed in Table 3. A composite map illustrating the complete migration routes of all tagged individuals is presented in Figure 4.


[14] Wild Birds Regulation Unit (WBRU), *Satellite Tagging of Song Thrushes* (Turdus philomelos) *and Eurasian Golden Plovers* (Pluvialis apricaria altifrons)*: Results from the 2022 – 2025 Research Period*, Final Report (2026). Available online: https://wbru.gov.mt/wp-content/uploads/2026/09/Final_Report_GP-ST-satellite-tagging-2022-2025.pdf.


Table 3. GPS tracking data for Golden Plovers, detailing migratory routes, wintering and breeding locations.


A note about the formatting: Deployment locations and dates are in italics; the last recorded location and date of transmission are in bold. Inset maps show the location of the satellite-tagged individuals during the breeding season.



Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**208612**	*Birżebbuġa, Malta* **Birżebbuġa, Malta**	*16/12/2022* **17/12/2022** (shot; tag returned to WBRU the following working day)	
Pluvialis apricaria 	**208581**	*Birżebbuġa, Malta* Birżebbuġa, Malta Filfla, Malta Lampedusa Chebba, Tunisia Sidi Bou Ali, Tunisia Enfidha, Tunisia Bouficha, Tunisia Errmal, Tunisia Errmal, Tunisia Hammamet, Tunisia Menzel Temime, Tunisia Menzel Temime, Tunisia Haouaria, Tunisia Nabeul, Tunisia Marettimo, Sicily Favignana, Sicily Randazzo, Sicily Rombiolo, Italy San Gregorio d'Ippona, Italy Campobasso, Italy Rače, Slovenia Studená, Czechia Horka u Staré Paky, Czechia Gmina Brzeźnica, Poland Gmina Gryfice, Poland Grästorp Municipality, Sweden Årjäng Municipality, Sweden Arvika Municipality, Sweden Arvika Municipality, Sweden Härjedalen, Sweden Falkvålen, Sweden **Falkvålen, Sweden**	*30/12/2022* 01/01/2023 02/01/2023 03/01/2023 04/01/2023 09/01/2023 12/01/2023 19/01/2023 20/01/2023 18/02/2023 22/02/2023 26/02/2023 09/03/2023 11/03/2023 13/03/2023 15/03/2023 15/03/2023 17/03/2023 18/03/2023 18/03/2023 20/03/2023 22/03/2023 24/03/2023 26/03/2023 28/03/2023 31/03/2023 01/04/2023 03/04/2023 08/04/2023 08/05/2023 10/05/2023 14/05/2023 **27/07/2023**	 




Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**208591**	*Birżebbuġa, Malta*	*07/02/2023*	 
		Birżebbuġa, Malta	07/02/2023	
		Lampedusa	09/02/2023	
		Mahdia, Tunisia	10/02/2023	
		Mahdia, Tunisia	18/02/2023	
		Moknine, Tunisia	20/02/2023	
		Kalaa Kebira, Tunisia	22/02/2023	
		Sebhika Kelbia, Tunisia	24/02/2023	
		Kondar, Tunisia	26/02/2023	
		Hammamet, Tunisia	28/02/2023	
		Bou Argoub, Tunisia	02/03/2023	
		Dar Chaabane El Fehri, Tunisia	06/03/2023	
		Korba, Tunisia	08/03/2023	
		Haouaria, Tunisia	10/03/2023	
		Trapani, Sicily	12/03/2023	
		Messina, Sicily	14/03/2023	
		Reggio Calabria, Italy	14/03/2023	
		Foggia, Italy	16/03/2023	
		Radonić, Croatia	16/03/2023	
		Središče ob Dravi, Slovenia	18/03/2023	
		Parndorf, Austria	20/03/2023	
		Ebenthal, Austria	22/03/2023	
		Čelechovice na Hané, Czechia	24/03/2023	
		Gmina Świdwin, Poland	26/03/2023	
		Hylte Municipality, Sweden	28/03/2023	
		Gislaved Municipality, Sweden	30/03/2023	
		Torsby Municipality, Sweden	01/04/2023	
		Malung, Sweden	10/04/2023	
		Malung, Sweden	08/05/2023	
		Strömsund Municipality, Sweden	10/05/2023	
		Sorsele Municipality, Sweden	12/05/2023	
		Sorsele Municipality, Sweden	**26/07/2023**	


Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**208617**	*Birżebbuġa, Malta*	*26/02/2023*	 
		Marsaskala, Malta	26/02/2023	
		Syracuse, Italy	26/02/2023	
		Crotone, Italy	26/02/2023	
		Lecce, Italy	26/02/2023	
		Ozrinići, Montenegro	28/02/2023	
		Novakovići, Montenegro	28/02/2023	
		Kaluđerovići, Serbia	02/03/2023	
		Hercegovačka Goleša, Serbia	02/03/2023	
		Zaovine, Serbia	04/03/2023	
		Tomnatic, Romania	06/03/2023	
		Tiszavasvári, Hungary	08/03/2023	
		Mezőzombor, Hungary	08/03/2023	
		Drochlin, Poland	10/03/2023	
		Koniecpol, Poland	10/03/2023	
		Dłużniewo Duże, Poland	12/03/2023	
		Makauskai, Lithuania	12/03/2023	
		Lebedžiūnai, Lithuania	20/03/2023	
		Krape, Latvia	22/03/2023	
		Kārļi, Latvia	24/03/2023	
		Naukšēni Parish, Latvia	26/03/2023	
		Sõrandu, Estonia	28/03/2023	
		Auküla, Estonia	30/03/2023	
		Ristijärvi, Finland	01/04/2023	
		Ristijärvi, Finland	14/04/2023	
		Hyrynsalmi, Finland	16/04/2023	
		Pielavesi, Finland	18/04/2023	
		Pielavesi, Finland	22/04/2023	
		Tervo, Finland	24/04/2023	
		Keitele, Finland	26/04/2023	
		Tervo, Finland	08/05/2023	
		Pudasjärvi, Finland	10/05/2023	
		Pudasjärvi, Finland	12/05/2023	
		Sodankylä, Finland	14/05/2023	
		Northern Lapland, Finland	16/05/2023	
		Northern Lapland, Finland	**29/07/2023**	





Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**208612** (re-used)	*Birżebbuġa, Malta*	*12/03/2023*	
		Birżebbuġa, Malta	21/03/2023	
		Mediterranean Sea (268km from eastern coast of Malta)	23/03/2023	
		Idrousa, Greece (10km NE from Andros Island)	25/03/2023	
		Beylik, Turkey	02/04/2023	
		Black Sea (33km from northern coast of Hatipler, Turkey)	05/04/2023	
		Temryuksky District, Russia	14/04/2023	
		Gorodovikovsky District, Russia	17/04/2023	
		Gorodovikovsky District, Russia	20/04/2023	
		Zavetinsky District, Russia	23/04/2023	
		Beloozerskii, Republic of Kalmykia, Russia	23/04/2023	
		Sarpinsky District, Republic of Kalmykia, Russia	05/05/2023	
		Dizhiembet, Zhanybek District, Kazakhstan	08/05/2023	
		Dizhiembet, Zhanybek District, Kazakhstan	17/05/2023	
		Zhiyenbat, Zhanybek District, Kazakhstan	20/05/2023	
		Zhiyenbat, Zhanybek District, Kazakhstan	23/05/2023	
		Grishenka, Denisov District, Kazakhstan	26/05/2023	
		Grishenka, Denisov District, Kazakhstan	29/05/2023	
		Kostanay District, Kazakhstan	01/06/2023	
		Evenkiysky District, Krasnoyarsk Krai (Siberia), Russia	12/06/2023	
		Evenkiysky District, Krasnoyarsk Krai (Siberia), Russia	11/08/2023	
		Krasnoyarsk Krai (Siberia), Russia	23/08/2023	
		Krasnoyarsk Krai (Siberia), Russia	**04/09/2023**	




Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**208614**	*Birżebbuġa, Malta* Birżebbuġa, Malta Balaoum, Tunisia Dar el Oussef, Tunisia Kondar, Tunisia Kondar, Tunisia El Alam, Tunisia Mazara del Vallo, Trapani, Italy Gmina Żychlin, Poland Färingsö, Sweden Abisko Östra, Sweden Abisko Östra, Sweden Abisko Östra, Sweden **Abisko Östra, Sweden**	*20/11/2023* 22/11/2023 23/11/2023 24/11/2023 19/12/2023 13/01/2024 07/02/2024 03/03/2024 28/03/2024 22/04/2024 16/05/2024 10/06/2024 05/07/2024 **30/07/2024**	
Pluvialis apricaria 	**208615**	*Birżebbuġa, Malta*	*21/11/2023 (10:40am)*	No GPS data
Pluvialis apricaria 	**208616**	*Birżebbuġa, Malta*	*21/11/2023 (02:50pm)*	No GPS data


Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**208618**	*Birżebbuġa, Malta*	*26/11/2023*	 
		Birżebbuġa, Malta	27/11/2023	
		Ech-Chahda Est, Tunisia	30/11/2023	
		Ouled Moulahem Ouest, Tunisia	03/12/2023	
		Sebkhet de Sidi El Hani, Tunisia	28/12/2023	
		Kroussia Ouest, Tunisia	22/01/2024	
		Kondar, Tunisia	16/02/2024	
		Damousse, Tunisia	12/03/2024	
		Falkenberg, Sweden	06/04/2024	
		Råde Municipality, Norway	01/05/2024	
		Steinsdalen, Norway	26/05/2024	
		Sjøvegan, Norway	20/06/2024	
		Sjøvegan, Norway	15/07/2024	
		Sjøvegan, Norway	**09/08/2024**	







Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**208619**	*Birżebbuġa, Malta*	*08/12/2023*	 
		Birżebbuġa, Malta	14/12/2023	
		Linosa, Italy	18/12/2023	
		Bouhlale El Ali Nord, Tunisia	21/12/2023	
		Ech-Chahda Est, Tunisia	14/01/2024	
		Sidi El Heni Sud, Tunisia	08/02/2024	
		Bechachma, Tunisia	04/03/2024	
		Lučenec, Slovakia	29/03/2024	
		Olaine Municipality, Latvia	22/04/2024	
		Kittilä, Finland	16/05/2024	
		Inari, Finland	11/06/2024	
		Inari, Finland	06/07/2024	
		Inari, Finland	**31/07/2024**	
Pluvialis apricaria 	**208620**	*Birżebbuġa, Malta*	*11/01/2024*	No GPS data
		Birżebbuġa, Malta	**11/01/2024** (shot; tag returned to WBRU)	
Pluvialis apricaria 	**208626**	*Żurrieq, Malta*	*12/01/2024*	No GPS data
		Żurrieq, Malta	**12/01/2024** (possibly shot; tag not returned to WBRU)	








Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**208629**	*Birżebbuġa, Malta* Birżebbuġa, Malta Mezraia, Tunisia Sabkhet Sijoumi Sud, Tunisia Budzyń, Poland Larvik Municipality, Norway Kvaløya, Tromsø Municipality, Norway Kvaløya, Tromsø Municipality, Norway Kvaløya, Tromsø Municipality, Norway **Kvaløya, Tromsø Municipality, Norway**	*12/01/2024* 19/01/2024 13/02/2024 09/03/2024 03/04/2024 28/04/2024 23/05/2024 17/06/2024 12/07/2024 **06/08/2024**	
Pluvialis apricaria 	**208621**	*Birżebbuġa, Malta* **Birżebbuġa, Malta**	*13/01/2024* **17/01/2024** (shot; tag returned to WBRU)	No GPS data. Tag 208621 was decommissioned in September 2024 due to battery failure.





Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**208624**	*Birżebbuġa, Malta*	*14/03/2024*	
		Birżebbuġa, Malta	22/03/2024	
		Kulisheve, Dnipropetrovsk Oblast, Ukraine	15/04/2024	
		Domachi, Lipetsk Oblast, Russia	11/05/2024	
			04/06/2024	
		Priuralsky District, Yamalo-Nenets Autonomous Okrug, Russia (Western Siberia)		
		Tazovsky District, Yamalo-Nenets Autonomous Okrug, Russia (Western Siberia)	24/07/2024	
		Tazovsky District, Yamalo-Nenets Autonomous Okrug, Russia (Western Siberia)	**19/08/2024**	
Pluvialis apricaria 	**208620** (re-used)	*Birżebbuġa, Malta*	*06/11/2024*	
		La Marsa, Tunisia	14/03/2025	
		Månslunda, Sweden	02/04/2025	
		Tolsjö, Sweden	26/04/2025	
		Staldvik, Norway	21/05/2025	
		Røyrvik Municipality, Norway	29/05/2025	
		Hattfjelldal Municipality Norway	30/05/2025	
		Kroken, Norway	31/05/2025	
		Kautokeino Municipality, Norway	15/06/2025	
		Porsanger Municipality, Norway	28/06/2025	
		Porsanger Municipality, Norway	29/06/2025	
		Porsanger Municipality, Norway	**30/06/2025**	


Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**208627**	*Birżebbuġa, Malta*	*14/11/2024*	 
		Hlavani, Odesa Oblast, Ukraine	02/04/2025	
		Kuygenkol, Kazakhstan	26/04/2025	
		Kulstan, Kazakhstan	21/05/2025	
		Zhanazhol, Kazakhstan	29/05/2025	
		Novopokrovka, Tomsk Oblast, Russia	30/05/2025	
		Argat-Yul, Tomsk Oblast, Russia	31/05/2025	
		Evenkiysky District Krasnoyarsk Krai, Russia	15/06/2025	
		Evenkiysky District Krasnoyarsk Krai, Russia	28/06/2025	
		Evenkiysky District Krasnoyarsk Krai, Russia	29/06/2025	
		Evenkiysky District Krasnoyarsk Krai, Russia	**30/06/2025**	
Pluvialis apricaria 	**208634**	*Żurrieq, Malta*	*24/11/2024*	 
		Salemi, Trapani, Italy	14/03/2025	
		Sárkeresztúr, Hungary	02/04/2025	
		Ageri, Järva County, Estonia	26/04/2025	
		Rovaniemi, Finland	21/05/2025	
		Pohjois-Lappi, Finland	29/05/2025	
		Pohjois-Lappi, Finland	30/05/2025	
		Pohjois-Lappi, Finland	31/05/2025	
		Nesseby Municipality, Varangerbotn, Norway	15/06/2025	
		Tana Municipality, Norway	28/06/2025	
		Tana Municipality, Norway	29/06/2025	
		Tana Municipality, Norway	**30/06/2025**	

Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**208622**	*Birżebbuġa, Malta*	*24/11/2024*	
		Campomarino, Campobasso, Italy	14/03/2025	
		Slampe, Tukums Municipality, Latvia	02/04/2025	
		Hälvati, Pärnu County, Estonia	26/04/2025	
		Pohjois-Lappi, Finland	21/05/2025	
		Pohjois-Lappi, Finland	29/05/2025	
		Pohjois-Lappi, Finland	30/05/2025	
		Pohjois-Lappi, Finland	31/05/2025	
		Inari, Finland	15/06/2025	
		Inari, Finland	28/06/2025	
		Inari, Finland	29/06/2025	
		Inari, Finland	**30/06/2025**	

Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**208623**	*Birżebbuġa, Malta*	*26/11/2024*	
		Trigortsi, Bulgaria	02/04/2025	
		Lugovoye, Kazakhstan	21/05/2025	
		Polovnikovka, Kazakhstan	29/05/2025	
		Tarsky District Omsk Oblast, Russia	30/05/2025	
		Kargasoksky District Tomsk Oblast, Russia	31/05/2025	
		Evenkiysky District Krasnoyarsk Krai, Russia	15/06/2025	
		Evenkiysky District Krasnoyarsk Krai, Russia	28/06/2025	
		Evenkiysky District Krasnoyarsk Krai, Russia	29/06/2025	
		Evenkiysky District Krasnoyarsk Krai, Russia	30/06/2025	
		Evenkiysky District Krasnoyarsk Krai, Russia	10/07/2025	
		Taymyrsky Dolgano-Nenetsky District Krasnoyarsk Krai, Russia	04/08/2025	
		Taymyrsky Dolgano-Nenetsky District Krasnoyarsk Krai, Russia	**16/08/2025**	
Pluvialis apricaria 	**208635**	*Birżebbuġa, Malta*	*26/11/2024*	No GPS data
Pluvialis apricaria 	**208636**	*Birżebbuġa, Malta*	*26/11/2024*	No GPS data
Pluvialis apricaria 	**208637**	*Żurrieq, Malta*	*01/12/2024*	
		Żurrieq, Malta	**26/04/2025** (possibly shot; tag not returned to WBRU)	

Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**209224**	*Żurrieq, Malta* **Żurrieq, Malta**	*06/12/2024* **26/04/2025** (possibly shot; tag not returned to WBRU)	
Pluvialis apricaria 	**209227**	*Żurrieq, Malta*	*10/12/2024*	No GPS data
Pluvialis apricaria 	**209225**	*Żurrieq, Malta* Darëzezë e Re, Albania Merișani, Romania **Tendrivske, Kherson Oblast, Ukraine**	*12/12/2024* 14/03/2025 02/04/2025 **26/04/2025**	
Pluvialis apricaria 	**209226**	*Żurrieq, Malta* Hanshir al Raqubah, Tunisia Kam'yans'ke, Odesa Oblast, Ukraine Kimer-Sayek-Kul', Kazakhstan Zhambyl District, Kazakhstan Zhambyl District, Kazakhstan Kargasoksky District Tomsk Oblast, Russia Alexandrovsky District Tomsk Oblast, Russia Taymyrsky Dolgano-Nenetsky District, Krasnoyarsk Krai, Russia Taymyrsky Dolgano-Nenetsky District, Krasnoyarsk Krai, Russia Taymyrsky Dolgano-Nenetsky District, Krasnoyarsk Krai, Russia **Taymyrsky Dolgano-Nenetsky District, Krasnoyarsk Krai, Russia**	*13/12/2024* 14/03/2025 02/04/2025 26/04/2025 21/05/2025 29/05/2025 30/05/2025 31/05/2025 15/06/2025 28/06/2025 29/06/2025 **30/06/2025**	 

Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**208625**	*Birżebbuġa, Malta*	*14/12/2024*	
		Piatra – Olt, Romania	14/03/2025	
		Heniches'kyi District Kherson Oblast, Ukraine	02/04/2025	
		Bychina, Perm Krai, Russia	21/05/2025	
Pluvialis apricaria 	**208630**	*Birżebbuġa, Malta*	*18/12/2024*	
		Birżebbuġa, Malta	**26/04/2025** (possibly shot; tag not returned to WBRU)	
Pluvialis apricaria 	**208631**	*Birżebbuġa, Malta*	*18/12/2024*	
		El Haouaria Sud, Tunisia	14/03/2025	
		Preajba de Jos, Romania	02/04/2025	
		Pysarivka, Luhansk Oblast, Ukraine	26/04/2025	
		Cherdynskiy Rayon, Perm Krai, Russia	21/05/2025	
		Kamgort, Perm Krai, Russia	29/05/2025	
		Beryozovsky District, Russia	30/05/2025	
		Nadymsky District, Russia	31/05/2025	
		Nadymsky District, Russia	15/06/2025	
		Tazovsky District, Russia	28/06/2025	
		Tazovsky District, Russia	29/06/2025	
		Tazovsky District, Russia	**30/06/2025**	
Pluvialis apricaria 	**208633**	*Birżebbuġa, Malta*	*16/01/2025*	
		Lapsi, l/o Siġġiewi, Malta	**30/05/2025** (shot; tag retrieved on 04/06/2025 by WBRU, assisted by EPU)	

Species	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Pluvialis apricaria 	**208628**	*Birżebbuġa, Malta*	*10/02/2025*	No GPS data
Pluvialis apricaria 	**208633** (re-used)	*Birżebbuġa, Malta*	*31/12/2025*	No GPS data




Figure 4: Composite map illustrating the wintering grounds, breeding grounds, and migration routes of satellite-tagged Golden Plovers (data from 30 Dec 2022 to 16 Aug 2025). Source: movebank.org, study name "Migration of Golden Plovers and Song Thrushes", study ID 1995912295.

Table 4 lists the last GPS locations recorded during the breeding season at the time this report was prepared.

Table 4. A summary of breeding grounds for satellite-tagged Golden Plovers.

Tag number	Last set of GPS fixes from overseas countries during the breeding season		Number of GPS fixes
	Breeding grounds	**Breeding period** (first–last fixes) (dd/mm/yyyy)	
208581	Falkvålen, Sweden	14/05/2023 – 27/07/2023	18
208591	Sorsele Municipality, Sweden	12/05/2023 – 26/07/2023	19
208617	Northern Lapland, Finland	16/05/2023 – 29/07/2023	19
208612	Evenkiysky District, (Central Siberia), Russia	12/06/2023 – 11/08/2023	5
208614	Abisko Östra, Sweden	16/05/2024 – 30/07/2024	4
208618	Sjøvegan, Norway	20/06/2024 – 09/08/2024	3
208619	Inari, Finland	11/06/2024 – 31/07/2024	3
208629	Kvaløya, Norway	23/05/2024 – 06/08/2024	4
208624	Tazovsky District (Western Siberia), Russia	24/07/2024 – 19/08/2024	2
208620	Porsanger Municipality, Norway	28/06/2025 – 30/06/2025	3
208627	Evenkiysky District (Central Siberia), Russia	15/06/2025 – 30/06/2025	4
208634	Tana Municipality, Norway	28/06/2025 – 30/06/2025	3
208622	Inari, Finland	15/06/2025 – 30/06/2025	4
208623	Evenkiysky–Taymyrsky Dolgano-Nenetsky Districts (Central Siberia), Russia	31/05/2025 – 16/08/2025	7
209225	*Inconclusive [Russia]*	N/A	N/A
209226	Taymyrsky Dolgano-Nenetsky District (Central Siberia), Russia	15/06/2025 – 30/06/2025	4
208625	Bychina, Perm Krai, Russia	21/05/2025 –	1
208631	Tazovsky District (Western Siberia), Russia	28/06/2025 – 30/06/2025	3

Source: movebank.org, study name “Migration of Golden Plovers and Song Thrushes”, study ID 1995912295.

Population size

The latest Article 12 data for the 2019–2024 reporting period specifies that the Northeast European breeding population of the Golden Plover (*P. a. altifrons*) within the EU territory (Finland and Sweden) has a **minimum breeding population of 172,790 pairs**. Table 5 outlines Malta’s Golden Plover EU reference population based on satellite tracking data alongside the 2019–2024 population trends.

Table 5. Malta's Golden Plover EU reference population based on satellite tracking data† and current population trend (2019–2024)

Country	% of EU Population	Min. Breeding Pairs	Weighted Share of Tracked Birds†	Malta's EU Reference Population (Min. Pairs)	Short-term Trend Direction	Short-term Trend Magnitude (%)			Long-term Trend Direction	Long-term Trend Magnitude (%)			Malta's EU Reference Population Short-term Trend			Malta's EU Reference Population Long-term Trend		
													Calculated Change in Breeding Pairs (Relative to Historical Baseline)					
						Min (Lower Bound)	**Max** (Upper Bound)	**Best Value**		**Min** (Lower Bound)	**Max** (Upper Bound)	**Best Value**	**Min** (Lower Bound Change)	**Max** (Upper Bound Change)	**Best Value**	**Min** (Lower Bound Change)	**Max** (Upper Bound Change)	**Best Value**
Finland	49.5%	90,790	50.0%	45,395	Decreasing	-30	-6	-19	Stable	-24	25	-2	-19,455	-2,898	-10,648	-14,335	9,079	-926
Sweden	49.1%	82,000	50.0%	41,000	Stable	N/A	N/A	0	Increasing	10	50	30	N/A	N/A	0	3,727	13,667	9,462
Total	**98.6%**	172,790	100%	**86,395**									**-19,455**	**-2,898**	**-10,648**	**-10,608**	**22,746**	**8,535**
									Overall Population Change				**-30.00%**	**-6.00%**	**-10.97%**	**-10.94%**	**35.74%**	**10.96%**
									Trend Direction * (Reference population)				**Decreasing** (>10% change)	**Stable** (<10% change)	**Decreasing** (>10% change)	**Stable** (<20% change since 1980)	**Increasing** (>20% change since 1980)	**Stable** (<20% change since 1980)

Data sources: European Environment Agency (2026). Birds Directive reporting (Art. 12): 2019-2024. Available at: https://nature-art12.eionet.europa.eu/article12/summary?period=4&subject=Pluvialis+apricaria&reported_name=.

† Movebank satellite tracking data (2022–2025): movebank.org, study name "Migration of Golden Plovers and Song Thrushes", study ID 1995912295. Data from 17 satellite-tagged Golden Plovers showed that all individuals wintered in Tunisia before migrating to breeding grounds in Russia (n = 7), Norway (n = 4), Finland (n = 3) and Sweden (n = 3). Overall, 35.3% of the tracked individuals bred within the EU.

* **Short-term Min (-30.00%) and Max (-6.00%):** Calculated on the reporting subset of Finland (45,395 pairs / 86,395 total pairs × 100 = 52.5% of the reference population). **Short-term Best Value (-10.97%):** Calculated on the full reference population of Finland & Sweden (100%), with Sweden reporting a stable 0% trend magnitude.

For the purpose of calculating the "small numbers" criterion in relation to the relevant "population concerned" (the reference population), Table 6 shows the **minimum** number of breeding pairs of *Pluvialis apricaria altifrons*, together with other relevant information required for calculating the 1% mortality threshold. According to the British Trust for Ornithology (BTO)[15], the typical lifespan of Golden Plover is four years and the annual mortality rate of adults is 27%, "based on a 50-year study in The Netherlands using colour-mark sightings" (Sandercock, 2003).

Table 6. Minimum breeding population and mortality rate – Malta's EU reference population

	Golden Plover (*Pluvialis apricaria altifrons*)	Source
Minimum breeding population – weighted number of min. pairs (see 5th column in Table 5)	**86,395**	Article 12 reports for the 2019–2024 reporting period (https://nature-art12.eionet.europa.eu/article12/summary?period=4&subject=Pluvialis+apricaria&reported_name=); Wetlands International (2009: 180); Movebank satellite tracking data (2022–2025): movebank.org, study name "Migration of Golden Plovers and Song Thrushes", study ID 1995912295.
Mortality rate juveniles	53.33%	Das Kompendium der Vögel Mitteleuropas (KVM) / Birds of the Western Palearctic (BWP)
Mortality rate adults	27%	BTO
Breeding rate (young per pair)	3.64	Golden Plover: European Management Plan-2009–2011, Technical Report-2009-034, Golden Plover.

Calculation of 1% mortality rate and "small numbers"

Golden Plover (*Pluvialis apricaria altifrons*)
Minimum breeding success: 86,395 x 3.64 = 314,478
Mortality rate of 1st year birds: 314,478 x 53.33% = 167,711
Mortality rate of adults: 172,790 x 27% = 46,653
Total annual mortality: 167,711 + 46,653 = 214,364
1% of total annual mortality: 214,364 x 1% = **2,144**

Autumn seasonal bag limit in relation to "small numbers"

Based on the 1% mortality rate (of the minimum population size), the "small numbers" calculation with respect to the national bag limit would amount to a figure below 2,144 Golden Plovers. However, it should be noted that the average number of Golden Plovers captured with clap-nets during period 2006–04 January 2026 is 484 (Table 2). Thus, the previous season's bag limit of 700 Golden Plovers should be retained if a derogation to open an autumn live-capturing season for this species is applied in 2026. This equates to 0.327% of the total annual mortality of the reference population (minimum population size), which is significantly less than the 1% threshold.

- **Golden Plover** – maximum national bag limit of **700** birds.

[15] Golden Plover *Pluvialis apricaria* Key Facts. British Trust for Ornithology. Available at: https://www.bto.org/learn/about-birds/birdfacts/golden-plover.

3. Conservation Status of Song Thrush (*Turdus philomelos*)

The Song Thrush (*Turdus philomelos*) is a passerine species in the thrush family found in woodland and forest ecosystems. It is a huntable species, listed under Annex II/B of the Birds Directive and is present throughout Europe and the northern part of Asia eastwards of Lake Baikal. **The northern populations are migratory and winter in western and southern Europe**. The European population is estimated at 24.4 – 38.4 million breeding pairs. Europe forms c.65% of the global range. According to BirdLife International (2026)[16], the global population size is estimated at 75 – (118)120 million individuals.

This species has an extremely large range (23,300,000 km²) and hence does not approach the thresholds for Vulnerable under the range size criterion (Extent of Occurrence <20,000 km^2 combined with a declining or fluctuating range size, habitat extent/quality, or population size and a small number of locations or severe fragmentation). The population trend appears to be increasing, and hence the species does not approach the thresholds for Vulnerable under the population trend criterion (>30% decline over ten years or three generations). The population size is extremely large and hence does not approach the thresholds for Vulnerable under the population size criterion (<10,000 mature individuals with a continuing decline estimated to be >10% in ten years or three generations, or with a specified population structure). For these reasons the species is evaluated as Least Concern (BirdLife International, 2026).

Global and European distribution

It has been documented that most Song Thrushes from the northern and eastern parts of the European range migrate via three or four main routes in a south-westerly direction (Ashmole M.J., 1962; Busse P & Maksalon L, 1986). With reference to the central Mediterranean region, the Italian publication *Atlante della Migrazione degli Uccelli in Italia*[17] states that "the Song Thrush is a regular breeder in Italy and a common passage migrant with the first thrushes occurring from late August, but high frequencies reached from late September, with the largest influxes in October with a maximum reached in the last decade of the month and good numbers till early November". With reference to the migratory pattern of this species, the authors note that:

"*Autumn recoveries mainly involve northern and central Italy, where most records are on the western side. Much less data from lower latitudes, while on islands most observations are from southern Sardinia. The good sample of direct autumn recoveries indicates NE–SW movements, primarily from the Balkans[18]. This main direction is similar to what [is] reported from more northern countries, like the UK, Norway, The Netherlands and Denmark. In our case Italy gets influxes from more eastern areas, with many birds entering through Friuli. Parallel routes towards our country are confirmed by the southernmost recoveries along the peninsula being referred to birds ringed in more eastern longitudes within the general ringing area.*

The national sample of first-capture data indicates a progressive decline in average wing length during the autumn, suggesting an earlier passage of more northern populations...The importance of the Mediterranean as a wintering area for the species is confirmed by the many recoveries, mainly concentrated south of the Apennines and with a concentration of data in Sardinia, although this island might be also affected by earliest return movements...Average size of birds ringed in Italy during the winter months show a further seasonal decline in size

[16] BirdLife International (2026) Species factsheet: *Turdus philomelos*. Downloaded from http://www.birdlife.org on 31/08/2026.
[17] Spina F. and Volponi S. (2008) *Atlante della Migrazione degli Uccelli in Italia*. Vol 2. Passeriformi (*Turdus philomelos*: pp.228–235).
[18] For the purpose of this conservation status report, the Balkans are limited to EU Member States: Bulgaria, Croatia, Greece, Romania and Slovenia.

till the central decade of December, followed by a new increase with the first decade of January, again suggesting early movements of birds of northern and eastern origin...Within a general framework of NE–SW Song Thrush movements across Europe, birds ringed in Italy move towards the coast of France, before reaching Mediterranean Iberia and North Africa, mainly in Algeria, later moving eastwards through Tunisia and flying back north via Sardinia and Corsica." Spina & Volponi (2008: 235).

Cramp *et al.* (1988)[19] note that the Song Thrush is mostly resident but northern populations are partially or entirely migratory and that in contrast to other species such as Redwing (*Turdus iliacus*) and Fieldfare (*Turdus pilaris*), populations show strong affinity to regular wintering areas. The authors further note that:

"*Most nominate* philomelos *from Fenno-Scandia, Germany, Switzerland, Poland, and [former] USSR are migratory, moving south-west or south-east through Europe to winter in southern England, France (mainly towards south-west), Spain and Portugal. Those from further north, especially first-year birds, winter furthest south to Canary Islands, Morocco, Algeria, Tunisia, Libya and Cyprus. Birds from Denmark, Netherlands, Belgium, and north-east France are partially resident with most others moving only short distances south or south-west, though considerable numbers from Netherlands winter in Britain and Ireland. Birds from Switzerland move south, with recoveries in southern France along Mediterranean coast into northern Italy, and in Corsica and Libya. Birds from east-central Europe winter correspondingly east of birds from Fenno-Scandia and western Europe: mainly in Italy, Yugoslavia, Greece, Balkans and Cyprus; also a very few recoveries in Balearics, Spain and Portugal.*

Populations from further east presumably winter largely from eastern Mediterranean to Iran...In early part of winter, migrant populations from Britain, Belgium, Netherlands and Denmark are chiefly in north-west France, north of Scandinavian birds (in southern Spain and Portugal), but by January birds from Low Countries have also moved south-west into Iberia. Southward departures in autumn begin in August but main passage September to early November. Birds from Fenno-Scandia move south-west on broad front, fringe of movement (or birds drifted west) passing through eastern Britain. Siberian birds depart mid-September while populations further south remain until November, in southern central Asia, Kazakhstan, and Turkmenistan; rarely recorded on passage outside breeding range. Birds wintering around Mediterranean arrive mid-October with frequent influxes until mid-April. During severe weather over Europe, large-scale mid-winter arrivals occur regularly in North Africa." Cramp *et al.* (1988: 990-992).

According to BirdLife International (2025) this species does not approach the thresholds for Vulnerable neither under the range size criterion, under the population trend criterion, nor under the population size criterion. For these reasons BirdLife International evaluates this species as Least Concern within both Europe and the EU[20] with an increasing population trend. IUCN[21] also classifies this species as **Least Concern**.

[19] Cramp S. *et al* (1988) Handbook of the Birds of Europe, the Middle East and North Africa: The Birds of the Western Palearctic. Vol V: Tyrant Flycatchers to Thrushes. The Royal Society for the Protection of Birds.
[20] http://www.birdlife.org/datazone/userfiles/file/Species/erlob/summarypdfs/22708822_turdus_philomelos.pdf
[21] http://www.iucnredlist.org/details/22708822/0

Article 12 reports (2019–2024)

This update includes the current population trends and indices of Malta's EU reference population as reported by Member States in their Article 12 report for period 2019–2024[22]. The Article 12 (2019–2024) dashboard containing Member State data files has recently been made public, accessible at https://nature-art12.eionet.europa.eu/article12/summary. However, the overall EU population status for each species is still pending, awaiting the subsequent release of national summaries, final datasets, and the "State of Nature" report by December 2026.

During the previous reporting cycle (2013–2018), the Song Thrush was reported as having a breeding population size of 12,200,000–20,500,000 pairs (previously reported at 12,700,000–21,900,000 pairs during the 2008–2012 reporting cycle) and a breeding range size of 3,100,000km² (previously reported at 3,520,000km² during the 2008–2012 reporting cycle).

The EU population status of the Song Thrush was assessed as **Secure**, because the species does not meet any of the IUCN Red List criteria for threatened or Near Threatened, or the criteria for Depleted or Declining (the EU population or range has not declined by 20% or more since 1980). Data on the wintering population trends for this species was not available at the time this report was prepared (August 2025).

Pan-European Common Birds Monitoring Scheme (2025 update)

The latest dataset available from the European Bird Census Council (EBCC, 2026)[23] at the time the conservation status of the Song Thrush (*Turdus philomelos*) was updated (August 2026), lists this species trend classification as **Moderate Increase,** defined as a "significant increase, but not significantly more than 5% per year" (Figure 5).

When compared with the previous EBCC update, the short-term [ten-year (2015–2024)] trend **decreased by 5%** [from +11% to +6%], whereas in the long-term trend (1980–2024) the population **decreased by 2%** (from +7% to +5%).[24]

[22] Available at: https://nature-art12.eionet.europa.eu/article12/summary.

[23] *Trends of common birds in Europe, 2025 update*. Available at: https://pecbms.info/trends-and-indicators/species-trends/species/turdus-philomelos/.

[24] Comparison between the previous and current EBCC updates is for information purposes only. The EBCC updates provide composite data (combined data of all contributing countries) on the latest population trends at pan-European level, not at EU level—it includes data pertaining to countries that are not part of Malta's EU reference population. Moreover, since the EBCC updates do not provide segregated data on population levels and trends for each contributing country, it is not possible to use EBCC's composite data, even if combined at EU level only, to update the assessment of the species at Malta's EU reference population level. The data at EU level that is segregated by country, and hence available to the Maltese authorities to carry out an assessment of the status of the species at reference population level, is the one published by the European Environment Agency as part of the Birds Directive Article 12 reporting obligations.




Figure 5. Long-term trend classification for *Turdus philomelos*. Source: EBCC (2025 update). Available at: https://pecbms.info/trends-and-indicators/species-trends/species/turdus-philomelos/.

List of Countries:

Austria	1998 - 2024
Belgium-Brussels	1992 - 2024
Belgium - Flanders	2007 - 2024
Belgium-Wallonia	1990 - 2024
Bulgaria	2005 - 2024
Czech Republic	1982 - 2024
Denmark	1976 - 2024
Estonia	1984 - 2024
Finland	1975 - 2024
France	1989 - 2024
Germany East	1991 - 2023
Germany West	1989 - 2023
Hungary	1999 - 2024
Italy	2000 - 2024
Latvia	2005 - 2024
Lithuania	1994 - 2024
Luxembourg	2010 - 2024
Netherlands	1984 - 2024
Norway	1996 - 2024
Poland	2000 - 2024
Republic of Ireland	1998 - 2024
Romania	2007 - 2024
Slovakia	2005 - 2024
Slovenia	2008 - 2024
Spain	1998 - 2024
Sweden	1975 - 2024
Switzerland	1999 - 2024
United Kingdom	1966 - 2024

Song Thrush distribution within the territory of the European Union (EU)

It has been documented that most of the Song Thrush populations from the northern and eastern parts of the European range migrate via three or four main routes in a south westerly direction (Ashmole M.J., 1962, Busse P & Maksalon L., 1986). The Song Thrush breeds throughout the territory of the European Union (EU), except Malta and Cyprus (EEA, 2020).

The Song Thrush does not breed in Malta (neither in the wild, nor in captivity). Table 7 below lists the live-capturing data for Song Thrush in Malta and Gozo from 2006 to 2025.

Table 7. Live-capturing data for Song Thrush (2006–2025)

Year	Amount Captured
2006	420
2007	10
2008	552
2009	1,199
2010	5,027
2011	2,069
2012	44
2013	135
2014	292
2015	248
2016	230
2017	183
2018	511
2019	722
2020	651
2021	1,006
2022	1,014
2023	1,286
2024	1,773
2025	1,366
Total	18,738
Mean (20-year average)	937

Data sources:
2006–2012: *Carnet de Chasse* data, Malta Environment and Planning Authority.
2013–2025: Game Reporting System, Wild Birds Regulation Unit.

Reference Population

The "small numbers" requirement, within the meaning of Article 9(1)(c) of the Birds Directive should be understood as "any sample of less than 1% of the total annual mortality of the population in question (average value) for those species which are not to be hunted and **a sample in the order of 1% for those species which may be hunted.** 'Population in question' is to be understood, with regard to migratory species, as the population of those regions from which come the main contingents passing through the region to which the derogation applies during its period of application". Given that the Song Thrush qualifies as a huntable species under the Birds Directive (MT: Annex IIB), the "*small numbers*" requirement is understood as "*a sample in the order of 1% of the total annual mortality of the population in question*".

The Song Thrush (*Turdus philomelos*) shows a strong affinity to regular wintering areas and a general framework of NE-SW movements across Europe, with birds from northern and eastern Europe migrating to wintering grounds in southern and western Europe (Milwright R.D.P., 2006[25]). The species' migratory routes and wintering fidelity are well-documented in ornithological literature. For example, a study by Spina & Volponi (2008) details the migratory patterns, confirming a general NE-SW migratory axis. The first two ring recoveries in Malta for this species were recorded in October 2024. The WBRU was notified of these recoveries by the Federazzjoni Kaċċaturi Nassaba Konservazzjonisti (FKNK), and subsequently, the ringing data was confirmed with the national ringing scheme. Both birds were ringed in Romania—one ringed as an adult on 16 April 2024, and the other ringed as a first-year on 2 October 2024, providing initial data on the species' presence and origin in the region.

[25] Milwright, R. D. P. (2006). The migration of European thrushes: a comparative study based on ringing recoveries. *Ringing & Migration*, 23(1), 1–13.

Based on this literature review, there is sufficient scientific evidence to determine that the EU Song Thrush populations migrating over Malta originate from the easternmost parts of the EU, specifically Romania, Bulgaria, Hungary, and Slovakia. This assertion is strongly corroborated by satellite tracking data[26], which offers precise information on migratory corridors and breeding grounds. The combination of these data sources supports the identification of the specific reference population for Malta (see Tables 10 and 11).

Satellite tracking data

As requested by the services of the European Commission in 2019, Malta initiated a research project to assess the migratory routes of this species using satellite tracking devices. The location of tagged specimens was monitored by the Maltese authorities using Argos satellite data repository to which a Movebank account was linked. The Movebank account showing all GPS fixes was converted to a publicly visible research study in 2021. This conservation status report incorporates the final findings of the study.

The study successfully tracked 21 Song Thrushes, with all birds wintering in Malta. Seven specimens provided conclusive evidence of their breeding locations in Romania (n=2), Bulgaria (n=2), Slovakia (n=1), Hungary (n=1), and Ukraine (n=1), confirming that 85.7% of the tracked Song Thrushes spent the breeding season within the European Union. The distinct SW-NE pre-nuptial migration route was also confirmed (Figure 6).

Wintering grounds (2022–2023 data only)

During Phase 1 and Phase 2 of the project, six Song Thrush tags out of a total of 14 deployed between 4 February 2022 and 10 March 2023 transmitted GPS fixes from continental Europe during pre-nuptial migration, after the specimens wintered in Malta (Table 8). The other tags did not transmit any data. The following table excludes data from the 2024 research period (Phase 3) since the seven specimens caught in 2024 were geotagged during the trailing end of the wintering period / start of post-nuptial migration (28 February – 3 March 2024).

Table 8. Song Thrush wintering period in Malta

Tag no.	Date tagged	Last GPS fix in Malta	First GPS fix overseas	Wintering period in Malta (days)		
				Minimum	Maximum	Mean
208582	04/02/2022	22/02/2022	29/03/2022	18	53	36
208583	04/02/2022	22/03/2022	12/04/2022	46	67	57
208590	06/12/2022	18/03/2023	06/04/2023	102	121	112
208593	09/12/2022	18/03/2023	07/04/2023	99	119	109
208578	23/01/2023	05/02/2023	07/04/2023	13	74	44
208585	28/02/2023	18/03/2023	06/04/2023	18	37	28
			Mean	**49**	**79**	**64**

The wintering location of the geotagged Song Thrushes was very restricted—all specimens remained in the same area where they were captured and satellite-tagged throughout the wintering period. As illustrated in Figure 6 further below, the flightpath taken by the geo-tagged Song Thrushes during pre-nuptial migration corroborates Spina and Volponi's (2008) observation of *a general framework of NE–SW Song Thrush movements across Europe*[27]*.*

Breeding grounds (2022–2024 data)

[26] Wild Birds Regulation Unit (WBRU), *Satellite Tagging of Song Thrushes* (Turdus philomelos) *and Eurasian Golden Plovers* (Pluvialis apricaria altifrons)*: Results from the 2022 – 2025 Research Period*, Final Report (2026). Available online: https://wbru.gov.mt/wp-content/uploads/2026/09/Final_Report_GP-ST-satellite-tagging-2022-2025.pdf.

[27] See p.12 of the Golden Plover and Song Thrush Conservation Status Report (2021 update). Available at: https://wbru.gov.mt/wp-content/uploads/2024/04/consStaRepMay21GpSt-1.pdf.

The Song Thrush fitted with tag no. **208582** remained in Romania throughout the breeding season (Table 9, first row) whereas the specimen fitted with tag no. **208583** appears to have remained in Belarus. However, since the last confirmed data was transmitted on 26 April 2022, it cannot be ascertained whether the specimen remained in Belarus throughout the breeding season or whether it migrated further north in May.

Deployments in 2023 (Phase 2) did not yield conclusive data on the location of the specimens' breeding grounds since the tags' batteries ran out after the first GPS fix in continental Europe (Table 9, rows 2–6). Deployments during the last phase of the Song Thrush geo-tagging project (Phase 3 [2024]) yielded comprehensive data on the breeding grounds of six of out the seven geotagged specimens (Table 9, rows 7–9 and 11–13).

Table 9. Song Thrush breeding territory



Tag number	Last set of GPS fixes from overseas countries during pre-nuptial and breeding period		Number of GPS fixes	Location of GPS fixes during breeding period
	Location	Date(s)		
208582	Romania	05/04/2022 – 14/06/2022	7	Romania
208583	Belarus	26/04/2022	1	*Inconclusive*
208590	Albania	06/04/2023	1	*Inconclusive*
208593	Greece	07/04/2023	1	*Inconclusive*
208578	Serbia	07/04/2023	1	*Inconclusive*
208585	North Macedonia	06/04/2023	1	*Inconclusive*
208598	Bulgaria	26/04/2024 – 01/06/2024	7	Bulgaria
208602	Bulgaria	26/04/2024 – 01/06/2024	7	Bulgaria
208605	Slovakia	08/05/2024 – 07/07/2024	9	Slovakia
208609	*No data*	–	–	–
208604	Hungary	26/04/2024 – 07/07/2024	10	Hungary
208600	Romania	26/03/2024 – 07/07/2024	8	Romania
208599	Ukraine	15/05/2024 – 07/07/2024	8	Ukraine




Table 10 below lists all Song Thrush GPS fixes recorded to date using Movebank online platform, including deployments from the previous reporting periods. Maps showing the migration route and last confirmed location (bold text) are also provided.



The GPS data from the deployments during Phase 1 and Phase 3 has shown that seven geo-tagged Song Thrushes bred in the following countries: Romania (n=2), Bulgaria (n=1), Slovakia (n=1), Hungary (n=1) and Ukraine (n=1) (refer to Table 11 further below on Malta's Song Thrush EU reference population).


Table 10. GPS tracking data for Song Thrushes, detailing migratory routes, wintering and breeding locations.



A note about the formatting: Deployment locations and dates are in italics; the last recorded location and date of transmission are in bold. Inset maps show the location of the satellite-tagged individuals during the breeding season.



Species	Ring No.	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Turdus philomelos	CC4804	**208582**	*Buskett, Malta*	*04/02/2022*	
			Buskett, Malta	08/02/2022	
			Buskett, Malta	15/02/2022	
			Buskett, Malta	22/02/2022	
			Planina, Serbia	29/03/2022	
			Ileanda, Romania	05/04/2022	
			Ileanda, Romania	12/04/2022	
			Ileanda, Romania	26/04/2022	
			Ileanda, Romania	03/05/2022	
			Ileanda, Romania	10/05/2022	
			Ileanda, Romania	17/05/2022	
			Ileanda, Romania	**14/06/2022**	
Turdus philomelos	CC4756	**208583**	*Buskett, Malta*	*04/02/2022*	
			Girgenti, Malta	08/02/2022	
			Girgenti, Malta	15/02/2022	
			Girgenti, Malta	22/02/2022	
			Buskett, Malta	22/03/2022	
			Baljci, Bosnia and Herzegovina	12/04/2022	
			Brgule, Bosnia and Herzegovina	19/04/2022	
			Kletsk District, Belarus	**26/04/2022**	
Turdus philomelos	CC4805	**208586**	*Girgenti, Malta*	*01/03/2022*	
			Girgenti, Malta	08/03/2022	
			Girgenti, Malta	15/03/2022	
			Girgenti, Malta	**22/03/2022**	


Species	Ring No.	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Turdus philomelos	CC4772	**208590**	*Buskett, Malta* Buskett, Malta Buskett, Malta Buskett, Malta Buskett, Malta **Likaj, Albania**	*06/12/2022* 26/12/2022 05/01/2023 16/01/2023 18/03/2023 **06/04/2023**	
Turdus philomelos	CC5303	**208589**	*Manikata, Malta*	*08/12/2022*	No GPS data
Turdus philomelos	CC4808	**208594**	*Girgenti, Malta* Girgenti, Malta Girgenti, Malta Girgenti, Malta Girgenti, Malta **Girgenti, Malta**	*09/12/2022* 26/12/2022 05/01/2023 16/01/2023 05/02/2023 **25/02/2023**	
Turdus philomelos	CC5605	**208593**	*Żebbiegħ, Malta* Mġarr, Malta Mġarr, Malta Mġarr, Malta Mġarr, Malta **Perama, Greece**	*09/12/2022* 26/12/2022 05/01/2023 16/01/2023 18/03/2023 **07/04/2023**	
Turdus philomelos	CC4773	**208595**	*Buskett, Malta*	*12/12/2022*	No GPS data

Species	Ring No.	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Turdus philomelos	CC4774	**208592**	*Buskett, Malta* Buskett, Malta Buskett, Malta **Buskett, Malta**	*12/12/2022* 26/12/2022 05/01/2023 **16/01/2023**	
Turdus philomelos	CC4776	**208597**	*Buskett, Malta* Buskett, Malta Buskett, Malta **Buskett, Malta**	*04/01/2023* 05/01/2023 16/01/2023 **04/02/2023**	
Turdus philomelos	CC4777	**208596**	*Buskett, Malta* Buskett, Malta Buskett, Malta Buskett, Malta Buskett, Malta Buskett, Malta **Buskett, Malta**	*16/01/2023* 04/02/2023 05/02/2023 25/02/2023 06/03/2023 18/03/2023 **06/04/2023**	
Turdus philomelos	CC4358	**208578**	*Buskett, Malta* Buskett, Malta Buskett, Malta **Bosnjane, Serbia**	*23/01/2023* 04/02/2023 05/02/2023 **07/04/2023**	

Species	Ring No.	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Turdus philomelos	CC3563	**208585**	*Simar, Malta* Simar, Malta Miżieb, Malta **Debreshte, North Macedonia**	*28/02/2023* 06/03/2023 18/03/2023 **06/04/2023**	
Turdus philomelos	CC5355	**208584**	*Wied Ħarq Ħamiem, Malta*	*10/03/2023*	No GPS data
Turdus philomelos	CC4815	**208598**	*Buskett, Malta* Buskett, Malta Buskett, Malta Buskett, Malta Kratovo, North Macedonia Chiprovtzi, Bulgaria Chiprovtzi, Bulgaria Chiprovtzi, Bulgaria Chiprovtzi, Bulgaria Chiprovtzi, Bulgaria Chiprovtzi, Bulgaria **Chiprovtzi, Bulgaria**	*28/02/2024* 29/03/2024 30/03/2024 31/03/2024 14/04/2024 26/04/2024 08/05/2024 15/05/2024 16/05/2024 17/05/2024 20/05/2024 **01/06/2024**	

Species	Ring No.	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Turdus philomelos	CC4816	**208602**	*Buskett, Malta*	*29/02/2024*	
			Buskett, Malta	03/03/2024	
			Buskett, Malta	26/03/2024	
			Buskett, Malta	02/04/2024	
			Debarca, North Macedonia	14/04/2024	
			Svoge, Bulgaria	26/04/2024	
			Svoge, Bulgaria	08/05/2024	
			Svoge, Bulgaria	15/05/2024	
			Svoge, Bulgaria	16/05/2024	
			Svoge, Bulgaria	17/05/2024	
			Svoge, Bulgaria	20/05/2024	
			Svoge, Bulgaria	**01/06/2024**	
Turdus philomelos	CC4817	**208605**	*Buskett, Malta*	*29/02/2024*	
			Buskett, Malta	29/03/2024	
			Buskett, Malta	30/03/2024	
			Buskett, Malta	31/03/2024	
			Buskett, Malta	02/04/2024	
			Rogatica Municipality, Bosnia and Herzegovina	14/04/2024	
			Békéssámson, Hungary	26/04/2024	
			Topoľa, Slovakia	08/05/2024	
			Topoľa, Slovakia	15/05/2024	
			Topoľa, Slovakia	16/05/2024	
			Topoľa, Slovakia	17/05/2024	
			Topoľa, Slovakia	20/05/2024	
			Topoľa, Slovakia	01/06/2024	
			Topoľa, Slovakia	13/06/2024	
			Topoľa, Slovakia	25/06/2024	
			Topoľa, Slovakia	**07/07/2024**	
Turdus philomelos	CC4818	**208609**	*Buskett, Malta*	*29/02/2024*	No GPS data

Species	Ring No.	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Turdus philomelos	CC4819	**208604**	*Buskett, Malta*	*03/03/2024*	
			Buskett, Malta	29/03/2024	
			Buskett, Malta	30/03/2024	
			Buskett, Malta	31/03/2024	
			Buskett, Malta	02/04/2024	
			Han Pijesak, Bosnia and Herzegovina	14/04/2024	
			Tállya, Hungary	26/04/2024	
			Nagyhuta, Hungary	08/05/2024	
			Kovácsvágás, Hungary	15/05/2024	
			Kovácsvágás, Hungary	16/05/2024	
			Kovácsvágás, Hungary	17/05/2024	
			Kovácsvágás, Hungary	20/05/2024	
			Kovácsvágás, Hungary	01/06/2024	
			Kovácsvágás, Hungary	13/06/2024	
			Kovácsvágás, Hungary	25/06/2024	
			Kovácsvágás, Hungary	**07/07/2024**	
Turdus philomelos	CC4820	**208600**	*Buskett, Malta*	*03/03/2024*	
			Bistra, Romania	26/03/2024	
			Posaga, Romania	14/04/2024	
			Posaga, Romania	17/05/2024	
			Posaga, Romania	20/05/2024	
			Posaga, Romania	01/06/2024	
			Posaga, Romania	13/06/2024	
			Posaga, Romania	25/06/2024	
			Posaga, Romania	**07/07/2024**	

Species	Ring No.	Tag No.	Location of GPS fixes	Date (dd/mm/yyyy)	Map
Turdus philomelos	CC4821	**208599**	*Buskett, Malta*	*03/03/2024*	
			Buskett, Malta	29/03/2024	
			Buskett, Malta	30/03/2024	
			Girgenti, Malta	02/04/2024	
			Litvytsya, Rivne Oblast, Ukraine	15/05/2024	
			Litvytsya, Rivne Oblast, Ukraine	16/05/2024	
			Litvytsya, Rivne Oblast, Ukraine	17/05/2024	
			Litvytsya, Rivne Oblast, Ukraine	20/05/2024	
			Litvytsya, Rivne Oblast, Ukraine	01/06/2024	
			Litvytsya, Rivne Oblast, Ukraine	13/06/2024	
			Litvytsya, Rivne Oblast, Ukraine	25/06/2024	
			Litvytsya, Rivne Oblast, Ukraine	**07/07/2024**	




Figure 6. Composite map of Song Thrushes pre-nuptial migration, illustrating general north-eastward movement (data from 04 Feb 2022 to 07 Jul 2024). Source: movebank.org, study name "Migration of Golden Plovers and Song Thrushes", study ID 1995912295.

Table 11. Malta's Song Thrush EU reference population based on satellite tracking data† and current population trend (2019–2024)

Country	% of EU Population	Min. Breeding Pairs	Weighted Share of Tracked Birds†	Malta's EU Reference Population (Min. Pairs)	Short-term Trend Direction	Short-term Trend Magnitude (%)			Long-term Trend Direction	Long-term Trend Magnitude (%)			Malta's EU Reference Population Short-term Trend			Malta's EU Reference Population Long-term Trend		
													Calculated Change in Breeding Pairs (Relative to Historical Baseline)					
						Min (Lower Bound)	**Max** (Upper Bound)	**Best Value**		**Min** (Lower Bound)	**Max** (Upper Bound)	**Best Value**	**Min** (Lower Bound Change)	**Max** (Upper Bound Change)	**Best Value**	**Min** (Lower Bound Change)	**Max** (Upper Bound Change)	**Best Value**
Bulgaria	2.3%	303,500	33.3%	101,167	Stable	-10	10	N/A	Increasing	N/A	N/A	82	-11,241	9,197	N/A	N/A	N/A	45,581
Hungary	2.1%	281,000	16.7%	46,833	Increasing	6	56	23.7	Unknown	N/A	N/A	N/A	2,651	16,812	8,973	N/A	N/A	N/A
Romania[1]	(5.2%)	1,585,519	33.3%	528,506	Increasing	46	120	83	Unknown	N/A	N/A	N/A	166,516	288,276	239,705	N/A	N/A	N/A
Slovakia	3.8%	350,000	16.7%	58,333	Increasing	5	10	N/A	Increasing	5	10	N/A	2,778	5,303	N/A	2,778	5,303	N/A
Total	**13.4%**	2,520,019	100%	**734,840**									**160,704**	**319,588**	**248,678**	**2,778**	**5,303**	**45,581**
												Overall Population Change	**27.99%**	**76.96%**	**76.13%**	**5.00%**	**10.00%**	**82.00%**
												Trend Direction * (Reference population)	**Increasing** (>10% change)	**Increasing** (>10% change)	**Increasing** (>10% change)	**Unknown / Incomplete data**	**Unknown / Incomplete data**	**Unknown / Incomplete data**

Data sources: European Environment Agency (2026). Birds Directive reporting (Art. 12): 2019-2024. Available at: https://nature-art12.eionet.europa.eu/article12/summary?period=4&subject=Turdus+philomelos&reported_name=.

[1] https://reportnet.europa.eu/public/dataflow/1438 (% MS [Romania] inferred).

† Movebank satellite tracking data (2022–2024): movebank.org, study name "Migration of Golden Plovers and Song Thrushes", study ID 1995912295. Data from 7 satellite-tagged Song Thrushes showed that all individuals wintered in Malta before migrating to breeding grounds in Bulgaria (n = 2), Romania (n = 2), Hungary (n = 1), Slovakia (n = 1) and Ukraine (n = 1). Overall, 85.7% of the tracked individuals bred within the EU.

* **Short-term Best Value (76.13%):** Calculated on the reporting subset of Hungary & Romania (575,339 pairs / 734,840 total pairs × 100 = 78.3% of the reference population). **Long-term Min (5.00%) and Max (10.00%):** Calculated on Slovakia only (58,333 pairs / 734,840 total pairs × 100 = 7.9%). **Long-term Best Value (82.00%):** Calculated on Bulgaria only (101,167 pairs / 734,840 total pairs × 100 = 13.8%).

Table 12 below shows the **minimum** number of Song Thrush breeding pairs, together with other relevant information required for the calculation of "small numbers". According to the British Trust for Ornithology (BTO)[28], the typical lifespan of Song Thrush is three years, and the annual mortality rate of adults is 43.7% ± 0.7%, which tallies with that specified by the *Birds of the Western Palaearctic (BWP)*.

Table 12. Minimum breeding population and mortality rate – Malta's EU reference population

	Song Thrush (*Turdus philomelos*)	Source
Minimum breeding population – weighted number of min. pairs (see 5th column in Table 11)	**734,840**	Article 12 reports for the 2019–2024 reporting period (https://nature-art12.eionet.europa.eu/article12/summary?period=4&subject=Turdus+philomelos&reported_name=); Cramp *et al* (1988); Spina & Volponi (2008); Movebank satellite tracking data (2022–2024): movebank.org, study name "Migration of Golden Plovers and Song Thrushes", study ID 1995912295.
Mortality rate – Juveniles	56%	Das Kompendium der Vögel Mitteleuropas (KVM) / Birds of the Western Palearctic (BWP)
Mortality rate – Adults	43.3%	BWP / British Trust for Ornithology (BTO)
Breeding rate (young per pair)	4	KVM / BWP / BTO

Calculation of 1% mortality rate and "small numbers"

Song Thrush (*Turdus philomelos*)
Minimum breeding success: 4 fledglings per pair (734,840 x 4) = 2,939,359
Mortality rate of 1st year birds: (2,939,359 x 56%) = 1,646,041
Mortality rate of adults: (1,469,679 x 43.3%) = 636,371
Total annual mortality: (1,646,041 + 636,371) = 2,282,412
1% of total annual mortality: (2,282,412 x 1%) = 22,824

Autumn seasonal bag limit in relation to "small numbers"

Based on the 1% mortality rate (of the minimum population size), the "small numbers" calculation with respect to the national bag limit would amount to a figure below 22,824 Song Thrushes. However, it should be noted that the average number of Song Thrushes captured with clap-nets during period 2006–2025 is 937 (Table 7). Thus, the previous season's bag limit of 5,000 Song Thrushes should be retained if a derogation to open an autumn live-capturing season for this species is applied in 2026. This equates to 0.219% of the total annual mortality of the reference population (minimum population size), which is significantly less than the 1% threshold.

- **Song Thrush** – maximum national bag limit of **5,000** birds.

[28] Song Thrush *Turdus philomelos* Key Facts. British Trust for Ornithology. Available at: https://app.bto.org/birdfacts/results/bob12000.htm.

4. Conclusion

Eurasian Golden Plover (*Pluvialis apricaria*)
The European (Eurasian) Golden Plover has an IUCN Least Concern classification. Since 2021, Estonia has been included in the "List of Countries" contributing to the latest European Bird Census Council (EBCC / PECBMS) updates, apart from Norway, Sweden and Finland (the latter two constituting Malta's EU reference population). According to the EU Management Plan for this species, Estonia does not form part of the Northeast European population of *Pluvialis apricaria altifrons* (Malta's EU reference population)[29].

Based on the latest dataset available from the European Bird Census Council (EBCC, 2025), there was a decrease in the combined breeding population of the contributing countries (Norway, Sweden, Finland and Estonia) both in the short-term and long-term trends. When compared with the previous EBCC update, the decrease in the short-term trend [ten-year slope] (2015–2024) was from -11% to -16% (an additional -5% decrease) and from -31% to -41% (an additional -10% decrease) in the long-term trend (1980–2024). The current EBCC assessment considers *Pluvialis apricaria* as Stable, defined as having "no significant increase or decline, and most probable trends are less than 5% per year".

Data from the current Birds Directive Article 12 report (2019–2024) show a short-term decreasing trend for Malta's EU reference population of *Pluvialis apricaria altifrons* (best value -10.97% based on combined Finnish and Swedish population data, with a minimum of -30% and maximum of +6% derived from Finnish data only). The species was reported as having a stable long-term trend (best value +10.96%, minimum -10.94%, maximum 35.74%).

Song Thrush (*Turdus philomelos*)
The Song Thrush has an IUCN Least Concern classification. The latest dataset available from the European Bird Census Council (EBCC, 2025) lists this species trend classification as Moderate Increase, defined as a "significant increase, but not significantly more than 5% per year".

When compared with the previous EBCC update, the short-term [ten-year (2015–2024)] trend decreased by 5% [from +11% to +6%], whereas in the long-term trend (1980–2024) the population decreased by 2% (from +7% to +5%).

Data from the current Birds Directive Article 12 report (2019–2024) show that Malta's reference population of *Turdus philomelos* (Bulgaria, Hungary, Romania and Slovakia) shows a short-term increasing trend (best value +76.13%, minimum +27.99%, maximum +76.96%), while its long-term trend is unknown due to missing data from three of the four constituent Member States.

[29] *The nominate southern subspecies* P. a. apricaria *nests in Ireland, Great Britain, Denmark, Germany, Latvia, Lithuania and Estonia. This subspecies winters in North-West Europe, form Ireland to South Britain, France and Iberia.* Source: EU Management Plan – Golden Plover (p.8).